

**Solutions to the “General Grand Unification Problem,”
and the Questions “How Did Our Universe Come Into Being?”
and “Of What is Empty Space Composed?”***

Robert A. Herrmann

Mathematics Department

U. S. Naval Academy

572 Holloway Rd.

Annapolis, MD 21402-5002

Presented before the Mathematics Association of America,
Western Maryland College, 12 NOV 1994.

Abstract: Using mathematical techniques to model one of the most simplistic of human linguistic processes, it is rationally predicted that within the nonstandard physical world there exists a force-like (logical) operator ${}^*\mathbf{S}$ and an entity w' such that ${}^*\mathbf{S}\{w'\}$ sequentially generates each of the Natural (i.e. physical) systems that comprise a Universe. This scientific model shows specifically that within the nonstandard physical world the behavior of each Natural world Natural-system is related logically. Further, the model predicts the rational existence of a single type of entity within the nonstandard physical world’s substratum that can be used to construct, by means of an exceptionally simple process, all of the fundamental Natural world particles used within particle physics. In important section 11.2, it is shown how (Natural law) allowable perturbations in Natural-system behavior are also included within this mathematical model. These results solve the pre-geometry problem of Wheeler. In general, the model predicts that when the behavior of these Universe creating processes are viewed globally, they can be interpreted as the behavior associated with a powerful intelligent agent.

CONTENTS

1. History	3
2. Two General Discussions	7
3. The Mathematics	15
3. Glossary	60
4. Two Important Explanatory E-mail Postings	64

*rah@usna.edu or drrahgid@hotmail.com

1. History.

[Note: References for this section appear on pages 4 – 5.] In August 1979, I was to give a paper before a gathering of mathematicians at the Summer Meeting of The American Mathematical Society [1]. This summer meeting is a joint meeting with the Mathematical Association of America and was being held at the University of Minnesota at Duluth. I boarded an airplane at Chicago for the last leg of my trip, and who should be sitting next to me at the window seat but John Wheeler, the Joseph Henry Professor of Physics emeritus at Princeton. Professor Wheeler was to deliver an invited talk before the Mathematical Association of America on what he believed was a fundamental difficulty with the philosophy of science associated with physical theories.

We discussed various things as we passed over some magnificent Minnesota thunderheads. Looking out of the window Professor Wheeler made the curious but simple statement, “We can’t do that.” What does this statement mean and who are the “we”?

After more discussion, the meaning of his comment was clear. First, the “we” are physicists and what they can’t do was illustrated by the thunderhead. The thunderhead is assumed to be a collection of Natural (i.e. physical) systems and, as a collection, can also display emerging properties. You have the water droplets, or crystals. Then their paths of motion caused by internal forces. The various electric potential differences produced by such motion and hundreds of other factors that scientists claim contribute to the overall behavior we observed from our window.

A Natural-system is a set or arrangement of physical entities that are so related or connected as to form an identifiable whole. Many Natural-systems have an associated physical theory. These theories are used to predict behavior for each Natural-system. BUT many of these theories use different methods to predict behavior. Indeed, one often has to restrict a particular theory to specific areas of application. When this is done you often have a boundary between two theories – an overlapping region – where one or the other but not both apply for the methods used in one of the theories are inconsistent with the methods used in the other theory. The hundreds of Natural-systems (now called subnatural-systems) that comprise the Natural system called a thunderhead are then put together in some manner to obtain the entire thunderhead one observes. With respect to the universe as a whole, the differences between various predicting theories are profound. What Professor Wheeler meant by the “. . . can’t do that” with respect to our universe is the acknowledgment that physicists had not found a general unifying theory that predicts all of the behavior of all subnatural systems that comprise the entire Natural-system called our universe. Science needed a theory that would be totally consistent, and a theory that would replace the piecemeal approach. Of course, such a unifying theory did not even exist for the behavior of the thunderhead we observed.

I restated this problem in the following manner.

Does Nature really combine subnatural-systems together, in a way that we can understand, to produce an entire Natural-system or does it use an entirely different method so that inconsistencies are somehow avoided? Is there something else required, something more basic than science has yet described and that’s needed to combine all the Natural-systems together that comprise our universe? Indeed, how can a universe that’s perceived by humans to have order and harmony really be a product of chaos?

By the way, I added that last question to this problem since it is certainly relevant.

After our discussion, it occurred to me that something discovered in October 1978, and now

called ultralogics [2], might possibly lead to an answer to these questions. But I also knew that the answer might be rather startling.

The standard approach is a piecemeal “bottom-up” approach beginning with the “bottom,” so to speak, of the scientific hierarchy (or “domain of explanation”). Supposedly, if you can find a unification of the fundamental forces (or interactions), then this would lead to an “upward” process that would eventually unify all physical theories. As discussed above, this is a doubtful assumption. Moreover, many scientists and philosophers of science believe that there are emergent properties of organisms that cannot be fully understood as products of DNA or chemistry and, thus, do not follow directly from the four fundamental forces.

On the other hand, there may be a *top-down* approach that answers the Wheeler questions. This means to find some scientific theory that gives processes which yield cosmologies, cosmologies that contain all of the processes that control the behavior of all Natural-systems.

The “deductive-world model,” (i.e. the D-world model) was the first constructed (1978) and is interpreted in a linguistic sense. D-world model properties will not be discussed directly within the following pages. Many of these properties can be found in sections 1 – 5 of the book *Ultralogics and More* [2]. However, some of what appears in the next sections does have the requisite linguistic interpretation associated with the D-world model. The actual model used to solve these problems is generally termed the *General Grand Unification Model (GGU-model)*. (For a special purpose, a portion of the GGU-model, the *metamorphic-anamorphosis model* (i.e. the MA-model) is employed.) Further, the intuitive concepts and mathematical methods that determine the GGU-model properties are very similar to those used for the additional D-world model conclusions. Thus becoming familiar with the GGU-model conclusions will aid in ones comprehension of the D-world model.

The basic construction of the GGU-model uses the following philosophical stance as described by deBroglie.

. . . *the structure of the material universe has something in common with the laws that govern the workings of the human mind* [3, p. 143]

It was determined that an appropriate cosmogony would depend upon the properties of a logic-like operator and subsidiary concepts associated with the *nonstandard physical world* (i.e. NSP-world) [4, part III]. The construction of the GGU-model began in August 1979 and a series of announcements [5 – 10] relative to its mathematical construction appeared in the Abstracts of papers presented to the American Mathematical Society. In particular, [5] mentions the logic-like operators.

In 1982, papers by Bastin [11], and Wheeler and Patton [12] were discovered in *The Encyclopedia of Ignorance*. I mention that [12] does not contain an important appendix that appears in the original 1975 paper. The paper by Bastin describes the *discreteness paradox* and the paper [13], using a few GGU-model procedures, presents a solution to this paradox. Next the requirements for an acceptable cosmogony stated in [12] were compared with GGU-model properties. A major difficulty in showing that the GGU-model meets all of these requirements is in the language used to give physical-like meaning to the abstract entities. After considerable reflection, the prefix “ultra” was decided upon. The use of this prefix is consistent with its use in the construction of the nonstandard structure. A relatively explicit construction of such a structure uses the concepts of the ultrafilter, the ultraproduct and the ultralimit. In 1987, a paper justifying the fundamental mathematical theory of nonstandard consequence operators was published [14].

Although a few technical aspects associated with the GGU-model were yet to be fully justified, two papers were written in 1986 [15], [16] announcing the solution to the general grand unification

problem and the pregeometry problem of Wheeler and Patton. To properly prepare paper [16], the original Wheeler and Patton paper [17] was utilized. An appendix in [17] describes how Wheeler and his colleagues at Princeton tried to construct a pregeometry from the statistics of very long propositions and very many propositions, where the term “proposition” refers to the propositional (i.e. the sentence) logical calculus. They failed to achieve a solution, but Wheeler left open the possibility that a solution could be obtained using concepts from the area of Mathematical Logic. This is exactly what has occurred.

Beginning in about 1989, a project was instituted to justify fully all of the GGU-model concepts, among other aspects of the NSP-world, and to present them in monograph form. The basic monograph [2] was completed in 1991 and the final result that completes the justification process was published in 1993 [18].

In the next section is given a very brief, general and mostly nonmathematical description of how the GGU-model and the language of the NSP-world solve the problems mentioned in the title. Presented in the third and technical section titled “The Mathematics,” with additional comments, is most of the actual mathematics that, when interpreted, describes the properties of the GGU-model [2] that correspond to the nonmathematical descriptions.

The following quotations are taken from [17] and are all relative to the Patton and Wheeler cosmogony requirements.

(1) *Five bits of evidence argue that geometry is as far from giving an understanding of space as elasticity is from giving a understanding of a solid.* [17, p. 539]

(2) *They also suggest that the basic structure is something deeper than geometry, that underlies both geometry and particles (“pregeometry”).* [17, p. 539]

(3) *For someday revealing this structure no perspective seems more promising than the view that it must provide the universe with a way to come into being.* [17, p. 539]

(4) *It brings us into closer confrontation than ever with the greatest questions on the book of physics: How did our universe come into being? And of what is it made?* [17, p. 540]

(5) *Tied to the paradox of the big bang and collapse is the question, what is the substance out of which the universe is made?* [17, p. 543]

(6) *But is it really imaginable that this deeper structure of physics should govern how the universe came into being? Is it not more reasonable to believe the converse, that the requirement that the universe should come into being governs the structure of physics?* [17, p. 558]

(7) *It is difficult to avoid the impression that every law of physics is “mutable” under conditions sufficiently extreme, . . .* [17, p. 568]

(8) *It is difficult to believe that we can uncover this pregeometry except as we come to understand at the same time the necessity of the quantum principle, with its “observer-participator,” in the construction of the world.* [17, p. 575]

(9) *. . . a guiding principle, is what we seek.* [17, p. 575]

As the GGU-model pregeometry is discussed, I will refer to these quotations where applicable. Relative to (9), the first half of the guiding principle is the deBroglie statement viewed from the NSP-world. The second half of the guiding principle is relative to the philosophy of realism and

an observation made by the originator of nonstandard analysis Abraham Robinson. First recall that Newton believed that infinitesimal measures were real measures associated with objectively real entities. Berkeley and Leibniz did not accept Newton's belief. Robinson at the end of his very first published paper on nonstandard analysis made the following statement relative to the modern concepts of how mathematical models are used to predict indirectly Natural-system behavior.

For phenomena on a different scale, such as are considered in Modern Physics, the dimensions of a particular body or process may not be observable directly. Accordingly, the question whether or not the scale of non-standard analysis is appropriate to the physical world really amounts to asking whether or not such a system provides a better explanation of certain observable phenomena than the standard system. . . . The possibility that this is the case should be borne in mind.

Fine Hall,
Princeton University [19, p. 440]

Robinson is referring to infinitesimal measures in this quotation. Since the publication of [19], nonstandard analysis has been applied to entities that are not infinitesimal in character. The second half of the guiding principle is an extension of this Robinson statement to the NSP-world. Thus the acceptance of the NSP-world as a viable realism depends upon whether or not it provides a "better" rational explanation of certain observable phenomena than the standard world model. The philosophy of what constitutes "a better explanation" is left to individual choice.

References

- [1] Herrmann, Robert. A. (1979). Perfect maps on convergence spaces I, *Notices of the AMS*, (26)(5): A-475.
- [2] Herrmann, Robert. A. (1991-93). The Theory of Ultralogics, See <http://www.serve.com/herrmann/bookss.htm> or <http://arxiv.org/abs/math/9903081> and <http://arxiv.org/abs/math/9903082>
- [3] deBroglie, L. (1963). In March, Arthur and Ira M. Freeman, *The New World of Physics*, Vintage Books, New York.
- [4] Herrmann, Robert A. (1989). Fractals and ultrasmooth microeffects, *J. Math. Physics*, 30(4) (April):805-808.
- [5] Herrmann, Robert A. (1981). Mathematical philosophy, *Abstracts of papers presented to the AMS*, Vol. 2(6). #81T-03-529:527.
- [6] Herrmann, Robert A. (1983). A useful *-real valued function, *Abstracts of papers presented to the AMS*, Vol. 4 (4), #83T-26-280:318.
- [7] Herrmann, Robert A. (1984). Nonstandard consequence operators I, *Abstracts of papers presented to the AMS*, Vol. 5 (1), #84T-03-61:129.
- [8] Herrmann, Robert A. (1984). Nonstandard consequence operators II, *Abstracts of papers presented to the AMS*, Vol. 5 (2), #84T-03-93:195.
- [9] Herrmann, Robert A. (1984). D-world alphabets I, *Abstracts of papers presented to the AMS*, Vol. 5 (4), #84T-03-320:269.
- [10] Herrmann, Robert A. (1984). D-world alphabets II, *Abstracts of papers presented to the AMS*, Vol. 5 (5), #84T-03-374:328.
- [11] Bastin, T., (1977). A clash of paradigms, in *The Encyclopaedia of Ignorance*, Duncan and Westen-Smith, eds. Pergamon, New York.

- [12] Wheeler, J. A. and C. M. Patton. (1977). Is physics legislated by cosmogony? in *The Encyclopaedia of Ignorance*, Duncan and Westen-Smith, eds. Pergamon, New York.
- [13] Herrmann, Robert A. (1983). Mathematical philosophical and developmental processes, *Nature and System*, 5(1/2):17–36.
- [14] Herrmann, Robert A. (1987). Nonstandard consequence operators, *Kobe J. Math.* 4(1):1–14.
- [15] Herrmann, Robert A. (1986). A solution to the grand unification problem, *Abstracts of papers presented to the AMS*, Vol. 7 (2), #86T-85-41:238.
- [16] Herrmann, Robert A. (1988) Physics is legislated by a cosmogony, *Speculations in Science and Technology*, 11(1):17–24.
- [17] Patton, C. M and J. A. Wheeler, (1975). Is physics legislated by cosmogony? in *Quantum Gravity: an Oxford Symposium*, Ishan, C., Penrose, R., and D. Sciama, eds, Clarendon, Oxford.
- [18] Herrmann, Robert A. (1993). A special isomorphism between superstructures, *Kobe J. Math.* 10(2):125-129.
- [19] Robinson, A., (1961). Non-standard analysis, *Nederl. Akad. Weimsch Proc.* Ser. A64 and *Indiag. Math.* 23:432 – 440.

Important Reference

The above reference [2], from which the mathematical portions of the monograph are taken, contains **all** of the fundamental concepts associated with the methods and processes used within the discipline of nonstandard analysis as they would apply to all aspects of the GGU-model and D-world model.

2. Two General Discussions

[Note: This first discussion is a portion of a general audience non-technical and elementary talk I gave on this subject. The numbers that appear in the double square brackets refer to the quotations from [17] as they appear on pages 5.] Let me point out that as long as scientists use mathematics to obtain their theories and written symbols, diagrams, photographs and the like to communicate their concepts then the GGU-model can't be eliminated. It will always be there lurking in the background.

Now to answer the question "How was our universe created?" Let's start by considering a single geometric point a few feet in front of you. A geometric point in this sense is a *position* in our universe and, for the present, has no other meaning. Now, I'm able to magnify this point for you by using a mathematical microscope with a power that's greater than any power that can ever be achieved by human means.

Suddenly, you see the point open up, like the iris of your eyes. What's revealed to you is a *background universe*, a *stratum*, or whatever you might like to call it. Now, the Natural-world is the world we can scientifically perceive, and this Natural-world point is still in your field of view with a small portion of the background universe surrounding it. You can't make out much detail, but there's definitely something there. The detail you see is sharper and clearer near to the single Natural-world point. Then clarity slowly fades as you proceed further from that one solitary Natural-world position within our universe. You can find no clear outer edge within your field of view. This background universe forms a portion of what I called the nonstandard physical world - the NSP-world ([4] above). [Note: The entire collection of all possible standard world points coupled with all of the NSP-world points that are infinitesimally near to them is called the *finite* or *bounded*

portion of the NSP-world.] The term the NSP-world is also used for other applications. However, for our purposes I'll discuss a portion of this NSP-world, the GGU-model.

Our universe is inside and “just as near to” the NSP-world as I have described it. One might say that this background universe is scientifically omnipresent. Now the GGU-model portion of this background universe specifically states that there will *never* be a human language that can give a completely detailed description for the mechanisms that may have produced our universe, but there do exist such mechanisms within this background universe. Such mechanisms exist but, no matter how hard we try, the human mind can't comprehend *all* of the *details*. Well then, what can be known about how universes, such as ours, can be created by NSP-world mechanisms? We can know general properties.

To begin with, there exists within the NSP-world a set of “things.” These “things” need not be considered as being within our Natural universe. But why do I call them “things”? I do know a lot about these “things.” But, unfortunately, if I were to give you a basic description for their contents, I would only use various mathematical terms from the fields of mathematical logic and set-theory. The most difficult task I've faced is to give some “physical” meaning to these “things.” These objects could easily fall into the category of those objects within the NSP-world that have NO human language physical descriptions at all. I know that these “things” behave like informational “superballs.”

After literally years of reflection, it was determined that these “things” can be described as containing all of the building plans, the laws of “Nature” and even step-by-step images of how an ideal universe will appear as it develops. They also contain a great deal of information that is incomprehensible to the human mind. It's interesting to note that, after I came to these conclusions, I came across a quotation from one of the greatest scientists of our time. Hermann Weyl is quoted as saying the following:

Is it conceivable that immaterial factors having the nature of images, ideas, 'building plans' also intervene in the evolution of the world as a whole?

We have “things” in the background universe that are ready, when the conditions are just right, to *aid* in the production of universes, ours included. And different “things” aid in the production of different universes. Actually, these “things,” which I have called *ultimate ultrawords*, don't work alone. Each ultraword is just one piece of an entire process. All pieces of the “puzzle” must be put together before a universe is produced. Now the existence of these ultrawords may not seem startling, but a lot more is yet to come.

What are the conditions that must exist in the NSP-world – conditions needed to trigger the creation of a universe such as ours? At present, there is no human scientific language that can detail these conditions, no human understanding of what these conditions are or were, only that such conditions exist.

I remind you that this cosmogony comes from a mathematical model, a model that can't be eliminated from modern science. This cosmogony satisfies all of the basic theoretical requirements of the scientific method. The universe in which we dwell, our solar system, the Earth, or a virus are Natural-systems. Natural-systems are studied by the physical scientist in a piecemeal fashion. They apply distinct procedures that seem to describe the moment-to- moment behavior of each distinct Natural-system. Now within the NSP-world there's one special process called “*S,” it's a hidden process, an *intrinsic* (hidden) *ultranatural process*. The *S-process is one of the entities know as

an *ultralogic*. When the conditions are just right, this force-like process takes one of these ultimate ultrawords and produces a universe. Indeed, it combines together, controls and coordinates all of the distinctly different Natural-systems that comprise a universe. The **S*-process applied to an ultimate ultraword produces each Natural-system and also yields the moment-to-moment alterations in the behavior of each and every Natural-system [2, 3, 4, 6]. But, how does it do this? We can't know many details, but a few simple mechanisms do present themselves.

Within the NSP-world there are objects, of a single type, (that I term *ultimate subparticles*) that, from the NSP-world viewpoint, can be “easily” combined together to produce every material object, electrons, protons, our earth, and everything else that might be termed material as well as immaterial fields, if such exist [2, 4, 5, 6]. This combining process cannot be reproduced by Natural means within any laboratory within our universe.

What about the development of our universe? That is how it changes with respect to time. This force-like process does produce a “beginning” for our universe and, as mentioned, in a remarkable step-by-step manner, it produces in the proper “time” ordered sequence all of the material changes from the very beginning until the universe arrives at a stage such as that which we observe about us. It produces all the Natural events that constitute the moment-to-moment changes that alter the appearance of a universe. This remarkable force-like process applied to an ultimate ultraword yields a solution to the general grand unification problem.

I say that this process is “remarkable” but I haven't explained why. I give you one example. From human perception, we often characterize certain Natural changes in a Natural-system as chaotic or random. This means that there seems to be no pattern for such changes – that is no pattern that can be comprehended by the human mind. Indeed, no human predictions can be made as to how individual objects will behave from one moment to another. From our viewpoint, there are no harmonious or regular laws that can produce such individual changes. It can be shown that from the NSP-world viewpoint the opposite is true. This seemingly irregular behavior within our universe is actually only what **we** can perceive of what is, in reality, an extremely regular process. How is this possible?

Well, it turns out that as a universe develops, as it changes, there are millions of other events taking place within the NSP-world that we can't perceive scientifically. These *ultranatural events* sustain and hold our universe together, so to speak. It's the ultranatural events that are combined together with the Natural events we perceive that actually comprise a complete change. Thus it's simply a matter of perception. I wish there were words in any language that fully describe this wondrous “combining” together process. There is a technical term, “ultrauniform,” that can be used. But this gives no indication of what is actually occurring. The steps in this combining together process are so minuscule, so small, so refined that the human mind can't fully appreciate nor comprehend the **S*-process. It doesn't correspond to anything that can at present be perceived or imagined by us.

Another startling aspect of the **S*-process is that every Natural event is connected, within the NSP-world, with every other Natural event. No Natural changes, from the NSP-world viewpoint, are independent from one another. A Natural event taking place in one galaxy is related by the **S*-process, to events taking place in every other galaxy. But, unfortunately, we can't know, except in general terms, the actual composition of an ultimate ultraword nor describe in any human language the necessary ultranatural events that sustain this process.

Now, what I've described, as best as I can, is the creation of an "ideal" universe. But what happens when there exist creatures within a universe that can alter its ideal development? Well, these creatures can only alter the Natural events, they can't alter the ultranatural events. No matter what these creatures do this "glue" that holds the universe together still remains [8]. I repeat, we can have no knowledge as to what these ultranatural events are. They cannot be described in a human language, ever. All we know is that they exist.

From the human perspective, there can be "sudden changes" in the behavior of a Natural-system at any time during its development [7]. But these changes are not truly "sudden" from the NSP-world viewpoint. Indeed, we can magnify a point in time, as we did with a point in space, and investigate what happens at the moment of change. Again wondrous events occur that are difficult to describe in a human language. Technically, the changes occur in an *ultrauniform* manner. This type of change, from the NSP-world viewpoint, can be compared with the mathematical concept of an "uniformly continuous" change. But ultrauniform is "infinitely" more uniform, so to speak, than the usual standard concept of an uniformly continuous change.

.
 If we look at the GGU-model as a whole, is there a way to describe it in its entirety? Yes, there is and this may be the most remarkable aspect of this research. The GGU-model can be characterized as behaving in its entirety like a super, super, super to an infinite degree, *mind* or possibly a *computer*. The processes are similar to how an almost inconceivably powerful mind or maybe a computer would behave.

In the next section titled "The Mathematics," is reproduced, with additional comments, some of the actual mathematics that, when interpreted, describes the properties of the GGU-model that correspond to the nonmathematical descriptions.

This portion of this discussion will be somewhat more technical than the previous portion. I will refer to section 3 as the motivation behind this mathematical model is discussed. The expression "Natural world" refers to the collection of all entities that are categorized as Natural-systems.

Relative to the behavior of a Natural-system, a general scientific approach is taken and it is assumed that scientists are interested in various types of descriptions for Natural-system behavior. It is not difficult to show that all forms of scientific description can be reduced to strings of symbols. Developmental paradigms are simply time related descriptions for the ideal behavior of any Natural-system viewed simply as strings of symbols. Although some of these collections of descriptions might be generated by a specific theory, a general approach that a developmental paradigm describes a sequence of Natural events is taken. The expression *Natural event* means an objectively real and physical occurrence that is categorized as "natural" by the physical scientist. This approach does not include any requirement that such a sequence be generated by some accepted and humanly comprehensible theory. On the other hand, theory generated sequences are not excluded.

On the first six pages of the next section are reproductions of pages taken from the paper "Nonstandard consequence operators" [Herrmann, R. A. *Kobe J. Math.* 4(1987): 1–14]. They detail how the basic somewhat unusual model $\mathcal{E}$ is constructed. The terminology used is that of the abstract mathematical structure. For example, certain subtle consequence operators are interpreted as ultralogics, while an ultraword is an unreadable sentence. The theorems on these pages show some of the behavior of nonstandard consequence operators. The most significant discussion on these first

six pages is the construction and embedding of the set $\mathcal{E}$. Although the results on these first six pages show some of the interesting behavior of nonstandard consequence operators they are not specifically needed to comprehend what follows. Further, the “time ordering” concept considered throughout what follows can be replaced with the notion of the “universal event number.”

Since an ultralogic is based upon the selection of some nontrivial logical process, there is a need to select a logical system that is common to all known logical systems used throughout scientific discourse. After some difficulty, a system, denoted by $\mathbf{S}$ and discussed at the beginning of section 7.3 titled ultrawords, was selected. I point out that the ordering of the sections of the included mathematics is not the same ordering in which the processes were originally discovered and used. Intuitively, throughout the modeling of these linguistic concepts, a specific “frozen segment” is considered as a description for a particular Natural event that occurs at a particular moment of time. From a scientific communication point of view, this description is all that can be scientifically known about such an event and is substituted for it.

Developmental paradigms, the deductive logic $\mathbf{S}$ and the like are embedded within a special but well-know mathematical structure called a superstructure by means of a fixed encoding and are further embedded into a nonstandard structure. The next step in the modeling process is to show that for a developmental paradigm written in a standard language there exists a new object, called an ultraword, that when the ultralogic (an intrinsic ultranatural process (i.e. an IUN-process)) $^*\mathbf{S}$ is applied to this ultraword the entire developmental paradigm (or the corresponding Natural event sequence) is logically produced. Theorem 7.3.1 establishes that for each developmental paradigm such an ultraword exists. Defining the *Natural world* as the collection of all Natural event sequences that correspond to the behavior of all systems that are categorized as Natural-systems, it follows that ultrawords cannot be entities within the Natural world under the GGU-model interpretation. Also the force-like operator $^*\mathbf{S}$ cannot be applied within the Natural world. Rather than simply accepting that ultrawords are “things” that cannot be further described, elsewhere additional intuitive meaning for this concept is discussed.

Relative to the collection of all events generated by the force-like process $^*\mathbf{S}$, a question that arises. Are only the developmental paradigm events obtained as a result of the process $^*\mathbf{S}$ applied to an ultraword w ? A conjecture was no. Theorem 7.3.2 shows that other “descriptions” for other types of events also occur when $^*\mathbf{S}$ is applied to a specific w . A later investigation, Theorem 10.1.1, shows the general composition of these new descriptions or events. But more importantly, Theorems 7.3.2 and 10.1.1 show that these new events must occur. Further, they cannot occur within the Natural world and cannot be described by any natural language. These events have been interpreted as events (called ultranatural events (i.e. UN-events)) that are needed to uphold and sustain a Natural-system’s development.

The above discussed results do not answered Wheeler’s basic question. The basic question is answered by Theorem 7.3.4 in an slightly more general mode. Every subnatural-system within the Natural-system called the universe can be associated with its own special system generating ultraword w'_i . Corollary 7.3.4.1 says that if we select a set of ultrawords that generate each and every subnatural-system that is within our Natural universe, then there (logically) exists an ultimate ultraword w' such that when the force-like process $^*\mathbf{S}$ is applied to w' each of the subnatural-system ultrawords are produced. Hence, all of the original subnatural-systems (the Natural event sequences) are produced by application of $^*\mathbf{S}$ to w' . Thus within the nonstandard physical world, not within the Natural world, there (logically) exists a force-like process the combines all of the Natural-systems

together in a consistent manner and yields our Natural universe and all of the moment-to-moment alterations that comprise its development. The mathematical existence of the ultimate ultraword w' yields a solution to the “general grand unification problem” as described by Wheeler.

Since the Natural event sequences are not necessarily predicted by a theory, then is it possible that the ideal developmental paradigms and the corresponding Natural event sequences are somehow “preselected”? Theorem 7.2.1 coupled with the discussion in 10.4 yields the necessary NSP-world IUN-selection processes. It is at this point that one of the most basic and significant properties of the GGU-model becomes apparent.

Suppose that you consider a partial denumerable developmental paradigm d_i where one of its members F_i is a frozen segment that represents a specific configuration for a Natural-system as it appears at a standard time interval $[t_i, t_{i+1})$. Notice that $\mathbf{F}_i \in \mathbf{T}_i$. The (cosmic or standard NSP-world) time t_i is conceived of as the moment of time in the past when a time fracture occurs and all other members of d_i represent behavior “after” such a time. There are infinitely many sequences (finite or denumerable) of standard frozen segments or *-frozen segments that can be adjoined to d_i , and that yield other developmental paradigms that describe Natural or ultranatural behavior for the same Natural-system but for cosmic times “prior to” t_i . This yields type d or type d' development paradigm.

Theorem 7.2.1 states that there is an external IUN-selection process that yields each of these developmental paradigms. A developmental paradigm that contains no additional frozen segments of any type prior to t_i along with its associated ultraword represents the maximum scenario. Note that a maximum scenario can yield by application of an ultralogic, *-frozen segments as well as frozen segments. A developmental paradigm that contains only time ordered standard frozen segments before and after t_i is a minimum scenario. An intermediate scenario contains some or all *-frozen segments representing ultranatural events prior to t_i . “Sudden alterations” are modeled by the minimum and intermediate scenarios and can be used for various purposes such as the Patton and Wheeler concept of “mutability” of Natural law or behavior. For the minimum and intermediate scenarios, can we investigate what happens during the NSP-world time when Natural law or Natural constants are altered? Can we “open up the time fracture,” so to speak, and look inside?

As discussed in section 7.5, applications of Theorem 7.5.1 yield a startling view of how these alterations are being made within the NSP-world. Furthermore, the force-like process $^*\mathbf{S}$ that produces all of the event sequences does so in a remarkable manner as described by Corollary 7.4.1.2.

As to various NSP-world objects that can mediate any GGU-model alterations, produce informational transmissions, and be considered as the composition of the vacuum, these are automatically generated by this mathematical model as shown by Theorem 9.3.1 where the descriptions are interpreted as descriptions for objects.

As to how the assumed “Laws of Nature” that seem to exist today came into being, the discussion in section 10.2 shows that these can also be assumed to have been produced by $^*\mathbf{S}$ applied to an ultraword. This answers another basic Wheeler question. The GGU-model cosmogony yields the various Natural laws that exist today. Further, an ultraword such as w gives an external unification to the collection of all written physical theories.

The last section in this paper deals with the “substance out of which the (Natural) universe is made.” It shows that using the method outlined in Theorem 9.3.1 the basic properties of subparticles

can be obtained. *Ultrasubparticles* are obtained with the aid of Theorem 11.1.1. Other properties of the infinitesimal and infinite hyperreal numbers coupled with a very simple hyperfinite translation (affine operation) lead to intermediate subparticles that can be finitely combined together to produce every basic entity within our Natural universe.

[Note (1): Relative to cyclic, multi-universe, plasma or any cosmology that claims that our universe has no Natural time beginning or no Natural time ending, such universes still have a “beginning” from the GGU-model viewpoint. No result in this paper is dependent upon a universe existing for only a finite period of time. Each Natural-system still has a beginning and ending with respect to an identified cycle. There is for each Natural-system within an i th cycle an ultraword w_j^i . Then there is a cycle ultimate ultraword $(w^i)'$. The multi-cycle-universe generating ultimate ultraword w' still exists and the force-like operator $*S$ still applies. That is, that w' generates each cycle and the contents of each cycle. Also note that if a Natural-system j is open-ended in that it continues to alter its appearance for all of time and has either no beginning or no ending in the intuitive sense, then this also can be modeled by considering a denumerable sequence of basic time intervals $[a_i, b_i)$. Again there is an ultraword w_j^i for each $[a_i, b_i)$ and, hence, an ultraword $(w_j)'$ that generates each of the w_j^i . Thus $(w_j)'$ generates the entire Natural-system's behavior. Finally, there is an ultimate ultraword w' that generates each $(w_j)'$ and, hence, each Natural-system. 3 MAR 1996]

[Note (2): For an ultimate GGU-model conclusion that shows how an GGU-model generated universe can be made to vary due to moment-to-moment perturbations, see section 11.2 that starts on page 55. This ultimate GGU-model conclusion also specifically models the “quantum” participatory requirements. 24 DEC 1996]

[Note (3): As mentioned previously, the set of all ultranatural initial conditions that would lead to universe creation can be entirely composed of entities that behave like “initial conditions” but these conditions cannot be described in any language associated with any entity within our universe. This fact does not mean that this set does not contain some ultranatural initial conditions that can be properly described by means of a comprehensible language. One such description follows the pattern of “field fluctuations.” The collection of all ultimate ideal ultrawords w' and its collection of perturbative ultimate ultrawords $\{w'_i\}$ within the NSP-world substratum can be supposed to form a dense collection W' . These collections fluctuate fortuitously with respect to NSP-world spacetime. Whenever a fluctuation exceeds a specified spacetime parameter, the ultralogic $*S$ acts upon the w' and a universe is generated. All of the universes that are generated by this process are disjoint. 2 JAN 1999]

[Note (4): Relative to the deBroglie statement and to all the areas that appear to be humanly comprehensible, with the exception of an additional postulate relative to the continuous energy spectrum for electromagnetic radiation that is required for subparticle generation, the GGU-model is based upon one and only one postulate. The postulate is that how nature combines together natural laws and processes to bring into being a natural event or a change in a natural system is modeled or mirrored by the mental processes human beings use to describe the produced events or changes. This hypothesis is verified by the largest amount of empirical evidence that could ever exist since whenever a scientist discusses or predicts natural system behavior human mental processes are applied. This includes all of the methods used to gather and analyze evidence that would tend to verify any scientific hypothesis. 3 JAN 1999]

In the next section, the references, when they have been reproduced, appear at the end of each

“chapter.”

Important Note

In the mathematics section 3, objects are chosen that seem to yield the simplest possible entities. This is done to minimize controversy and to allow most conclusions to be established in a convincing and straightforward manner. The philosophy of science employed is the exact same philosophy of science used in theoretical cosmology and quantum logic investigations. In almost all cases, the physical-like interpretations correlate directly to the mathematical structure. The first section of part 3 is a reproduction of a Kobe Math. J. paper. The second section of part 3 is a reproduction of portions of the book *Ultralogics and More* (see reference [2] of part 1) where all the details for the actual construction of the nonstandard model can be found. It should be self-evident that the results contained within this monograph are only the must basic and that further in-depth investigations should be pursued.

Intuitively, I am confident that all of the theorems are correct. If any “proof” is not convincingly established, then this should be easy to rectify. Finally, I do not contend that this is THE solution to this problem and questions. Although these results are speculative in character, they are no more hypothetical than the Everett-Wheeler-Graham many-worlds interpretation or Hartle-Hawking quantum gravity model. I do contend that since the method was devised in 1979 that these results are the FIRST such solutions obtained scientifically. The term “scientifically” refers to the sciences of concrete mathematical modeling, mathematical logic and interpreting mathematical structures physically.

3. The Mathematics

The following is a direct copy, with numerous many printers errors corrected, of the indicated portions of the indicated published paper.

Herrmann, R. A.

Kobe J. Math.

4(1987) 1-14

NONSTANDARD CONSEQUENCE OPERATORS

By Robert A. Herrmann

(Dedicated to Professor K. Iséki)

(Receiver October 22, 1984)

1. Introduction

In 1963, Abraham Robinson applied his newly discovered nonstandard analysis to formal first-order languages and developed a nonstandard logic [9] relative to the “truth” concept and structures. Since that time not a great deal of fundamental research has been attempted in this specific area with one notable exception [2]. However, when results from this discipline are utilized they have yielded some highly significant and important developments such as those obtained by Henson [3].

The major purpose for this present investigation is to institute formally a more general study than previously pursued. In particular, we study nonstandard logics relative to consequence operators [1] [5] [10] [11] defined on a nonstandard language. Since the languages considered are not obtained by the usual constructive methods, then this will necessitate the construction of an entirely new foundation distinctly different from Robinson’s basic embedding techniques. Some very basic results of this research were very briefly announced in a previous report [5].

In 2, we give the basic definitions, notations and certain new standard results are obtained that indicate the unusual behavior of the algebra of all consequence operators defined on a set. In 3, some new standard properties relative to subalgebras and chains in the set of all consequence operators are investigated. Finally, the entire last section is devoted to the foundations of the theory of nonstandard consequence operators defined on a nonstandard language.

2. Basic concepts

Our notations and definition for the standard theory of consequence operators are taken from references [1][5][10][11], and we now recall the most pertinent of these. Let L be any nonempty set that is often called a (sentential) *language*, $\mathcal{P}(L)$ denote the power set of L and for any set X let $F(X)$ denote the finite power set of X (i.e. the set of all finite subsets of X .)

DEFINITION 2.1 A mapping $C: \mathcal{P}(L) \rightarrow \mathcal{P}(L)$ is a consequence operator (or closure operator) if for each $X, Y \in \mathcal{P}(L)$

- (i) $X \subset C(X) = C(C(X)) \subset L$ and if

(ii) $X \subset Y$, then $C(X) \subset C(Y)$.

A consequence operator C defined on L is said to be *finite* (*finitary*, or *algebraic*) if it satisfies

(iii) $C(X) = \cup\{C(A) \mid A \in F(X)\}$.

REMARK 2.2 The above axioms (i) (ii) (iii) are not independent. Indeed, (i)(iii) imply (ii).

Throughout this entire article the symbol “ C ” with or without subscripts or with or without symbols juxtapositioned to the right will always denote a consequence operator. The only other symbols that will denote consequence operators are “ I ” and “ U ”. The symbol $\mathcal{C}$ [resp. $\mathcal{C}_f$] denotes the set of all consequence operators [resp. finite consequence operators] defined on $\mathcal{P}(L)$.

DEFINITION 2.3. (i) Let I denote the identity map defined on $\mathcal{P}(L)$.

(ii) Let $U: \mathcal{P}(L) \rightarrow \mathcal{P}(L)$ be defined as follows: for each $X \in \mathcal{P}(L)$, $U(X) = L$.

(iii) For each $C_1, C_2 \in \mathcal{C}$, define $C_1 \leq C_2$ iff $C_1(X) \subset C_2(X)$ for each $X \in \mathcal{P}(L)$. (Note that $\leq$ is obviously a partial order defined on $\mathcal{C}$.)

(iv) For each $C_1, C_2 \in \mathcal{C}$, define $C_1 \vee C_2: \mathcal{P}(L) \rightarrow \mathcal{P}(L)$ as follows: for each $X \in \mathcal{P}(L)$, $(C_1 \vee C_2)(X) = C_1(X) \cup C_2(X)$.

(v) For each $C_1, C_2 \in \mathcal{C}$, define $C_1 \wedge C_2: \mathcal{P}(L) \rightarrow \mathcal{P}(L)$ as follows: for each $X \in \mathcal{P}(L)$, $(C_1 \wedge C_2)(X) = C_1(X) \cap C_2(X)$.

(vi) For each $C_1, C_2 \in \mathcal{C}$ define $C_1 \vee_w C_2: \mathcal{P}(L) \rightarrow \mathcal{P}(L)$ as follows: for each $X \in \mathcal{P}(L)$, $(C_1 \vee_w C_2)(X) = \cap\{Y \mid X \subset Y \subset L \text{ and } Y = C_1(Y) = C_2(Y)\}$.

Prior to defining certain special consequence operators notice that $I, U \in \mathcal{C}_f$ and that I [resp. U] is a lower [resp. upper] unit for the algebras $\langle \mathcal{C}, \leq \rangle$ and $\langle \mathcal{C}_f, \leq \rangle$.

.....

4. Nonstandard Consequence Operators

Let $\mathcal{A}$ be a nonempty finite set of symbols. It is often convenient to assume that $\mathcal{A}$ contains a symbol that represents a blank space. As usual any nonempty finite string of symbols from $\mathcal{A}$, with repetitions, is called a *word* [8, p.222]. A word is also said to be an (intuitive) *readable sentence* [4, p. 1]. We let W be the intuitive set of all words created from the *alphabet* $\mathcal{A}$. Note that in distinction to the usual approach, W does not contain a symbol for the empty word.

We accept the concept delineated by Markov [3], the so-called “abstraction of identity,” and say that $w_1, w_2 \in W$ are “equal” if they are composed of the same symbols written in the same intuitive order (left to right). The *join* or juxtaposition operator between $w_1, w_2 \in W$ is the concept that yields the string w_1w_2 or w_2w_1 . Thus W is closed under join. Notice that we may consider a denumerable formal language as a subset of W .

Since W is denumerable, then there exists an injection $i: W \rightarrow \mathbb{N}$, where $\mathbb{N}$ is the set of natural numbers with zero. Obviously, if we are working with a formal

language that is a subset of W , then we may require i restricted to a formal language to be a Gödel numbering. Due to the join operator, a fixed member of W that contains two or more distinct symbols can be represented by various *subwords* that are joined together to yield the given fixed word. The word “mathematics” is generated by the join of $w_1 = \text{math}$, $w_2 = \text{e}$, $w_3 = \text{mat}$, $w_4 = \text{ics}$. This word can also be formed by joining together 11 not necessarily distinct members of W .

Let $i[W] = T$ and for each $n \in \mathbb{N}$, let $T^n = T^{[0,n]}$ denote the set of all mappings from $[0, n]$ into T . Each element of T^n is called a *partial sequence*, even though this definition is a slight restriction of the usual one that appears in the literature. Let $f \in T^n, n > 0$. Then the *order induced by f* is the simple inverse order determined by f applied to the simple order on $[0, n]$. Formally, for each $f(j), f(k) \in f[[0, n]]$, define $f(k) \leq_f f(j)$ iff $j \leq k$, where $\leq$ is the simple order for $\mathbb{N}$ restricted to $[0, n]$. In general, we will not use this notation $\leq_f$ but rather we will indicate this (finite) order in the usual acceptable manner by writing the symbols $f(n), f(n-1), \dots, f(0)$ from left to right. Thus we symbolically let $f(n) \leq_f f(n-1) \leq_f \dots \leq_f f(0) = f(n)f(n-1) \dots f(0)$.

Let $f \in T^n$. Define $w_f \in W$ as follows: $w_f = (i^{-1}(f(n)))(i^{-1}(f(n-1))) \dots (i^{-1}(f(0)))$, where the operator indicated by juxtaposition is the join. We now define a relation on $P = \cup\{T^n \mid n \in \mathbb{N}\}$ as follows: let $f, g \in P$. Then for $f \in T^n$ and $g \in T^m$, define $f \sim g$ iff $(i^{-1}(f(n))) \dots (i^{-1}(f(0))) = (i^{-1}(g(m))) \dots (i^{-1}(g(0)))$. It is obvious that $\sim$ is an equivalence relation on P . For each $f \in P$, $[f]$ denotes the equivalence class under $\sim$ that contains f . Finally, let $\mathcal{E} = \{[f] \mid f \in P\}$. Observe that for each $[f] \in \mathcal{E}$ there exist $f_0, f_m \in [f]$ such that $f_0 \in T^0$, $f_m \in T^m$ and if there exists some $k \in \mathbb{N}$ such that $0 < k < m$, then there exists some $g_k \in [f]$ such that $g_k \in T^k$ and if $j \in \mathbb{N}$ and $j > m$, then there does not exist $g_j \in T^j$ such that $g_j \in [f]$. If we define the *size* of a word $w \in W$ ($\text{size}(w)$) to be the number of not necessarily distinct symbols counting left to right that appear in W , then the $\text{size}(w) = m + 1$. For each $w \in W$, there is $f_0 \in T^0$ such that $w = i^{-1}(f_0(0))$ and such an $f_m \in [f_0]$ such that $\text{size}(w) = m + 1$. On the other hand, given $f \in P$, then there is a $g_0 \in [f]$ such that $(i^{-1}(g_0(0))) \in W$. Of course, each $g \in [f]$ is interpreted to be the word $(i^{-1}(g(k))) \dots (i^{-1}(g(0)))$.

Each $[f] \in \mathcal{E}$ is said to be a (formal) word or (formal) *readable sentence*. All the intuitive concepts, definitions and results relative to consequence operators defined for $A \in \mathcal{P}(W)$ are now passed to $\mathcal{P}(\mathcal{E})$ by means of the quotient map θ generated by the equivalence relation $\sim$. In the usual manner, the quotient map is extended to subsets of each $A \in \mathcal{P}(W)$, n -ary relations and the like. For example, let $w \in A \in \mathcal{P}(W)$. Then there exists $f_w \in P$ such that $f_w \in T^0$ and $f_w(0) = i(w)$. Then $\theta(i(w)) = [f_w]$. In order to simplify notation, the images of the extended (θi) composition will often be indicated by bold notation with the exception of customary relation symbols which will be understood relative to the context. For example, if T is a subset of W , then we write $\theta(i[T]) = \mathbf{T}$.

Let $\mathcal{N}$ be a superstructure constructed from the set $\mathbb{N}$ as its set of atoms. Our standard structure is $\mathcal{M} = (\mathcal{N}, \in, =)$. Let $^*\mathcal{M} = (^*\mathcal{N}, \in, =)$ be a nonstandard and elementary extension of $\mathcal{M}$. Further, $^*\mathcal{M}$ is an enlargement.

For an alphabet $\mathcal{A}$, there exists $[g] \in {}^*\mathcal{E} - \mathcal{E}$ such that there are only finitely many standard members of $\mathbb{N}$ in the range of g and these standard members injectively correspond to alphabet symbols in $\mathcal{A}$ [4, p. 24]. On the other hand, there exist $[g'] \in {}^*\mathcal{E} - \mathcal{E}$ such that the range of $[g']$ does not correspond in this manner to elements in $\mathcal{A}$ [4, p. 90].

Let $C \in \mathcal{H}$ map a family of sets $\mathcal{B}$ into $\mathcal{B}_0$. If C satisfies either the Tarski axioms (i), (ii) or (i), (iii), or the $*$ -transfer ${}^*(i)$, ${}^*(ii)$, or ${}^*(i)$, ${}^*(iii)$ of these axioms, then C is called a *subtle consequence operator*. For example, if $C \in \mathcal{C}$, then it is immediate that ${}^*\mathbf{C}: {}^*(\mathcal{P}(\theta(A)) \rightarrow {}^*\mathcal{P}(\theta(A)))$ satisfies ${}^*(i)$ and ${}^*(ii)$ for the family of all internal subsets of ${}^*(\theta(A))$. This ${}^*\mathbf{C}$ is a subtle consequence operator. For any set $A \in \mathcal{N}$, let ${}^\sigma A = \{ {}^*a \mid a \in A \}$. (In general, this definition does not correspond to that used by other authors.) If for a subtle consequence operator C there does not exist some similarly defined $D \in \mathcal{N}$ such that $C = {}^\sigma \mathbf{D}$ or $C = {}^*\mathbf{D}$, then C is called a *purely subtle consequence operator*. Let infinite $A \subset \mathcal{E}$ and $B = {}^*A - {}^\sigma A$. Then the identity $I: \mathcal{P}(B) - \mathcal{P}(B)$ is a purely subtle consequence operator.

There are certain technical procedures associated with the σ map that take on a specific significance for consequence operators. Recall that $\mathcal{N}$ is closed under finitely many power set or finite power set iterations. Let $X, Y \in \mathcal{N}$. It is not difficult to show that if $\mathcal{P}: \mathcal{P}(X) \rightarrow Y$, then for each $A \in \mathcal{P}(X)$, ${}^*(\mathcal{P}(A)) = {}^*\mathcal{P}({}^*A)$. Moreover, if $F: \mathcal{P}(X) \rightarrow Y$, where F is the finite power set operator, then for each $A \in \mathcal{P}(A)$, ${}^*(F(A)) = {}^*F({}^*A)$. If $C \in \mathcal{C}$ and $X \subset \mathcal{E}$, then $\mathbf{C}: \mathcal{P}(X) \rightarrow \mathcal{P}(X)$ has the property that for each $A \in \mathcal{P}(X)$, ${}^*(\mathbf{C}(A)) = {}^*\mathbf{C}({}^*A)$.

Recall that we identify each ${}^*n \in {}^*\mathbb{N}$ with $n \in \mathbb{N}$ since *n is but a constant sequence with the value n . Utilizing this fact, we have the following straightforward lemma the proof of which is omitted.

LEMMA 4.1. *Let $A \in \mathcal{N}$ and $C \in \mathcal{C}$.*

- (i) ${}^\sigma(F(A)) = F({}^\sigma A)$ and if $A \subset \mathcal{E}$, then ${}^\sigma(F(A)) = F(A)$.
- (ii) ${}^\sigma(\mathbf{C}(B)) = \mathbf{C}(B)$, for each $B \in \mathcal{P}(X)$, $X \subset \mathcal{E}$, and
- (iii) ${}^*\mathbf{C}[\{ {}^*A \mid A \in \mathcal{P}(X) \}] = \{ ({}^*A, {}^*B) \mid (A, B) \in \mathbf{C} \} = {}^\sigma \mathbf{C}$. If $F \in F(B)$, then ${}^\sigma(\mathbf{C}(F)) \subset ({}^\sigma \mathbf{C})({}^\sigma F) = ({}^\sigma \mathbf{C})(F)$. Also, ${}^\sigma(\mathbf{C}(B)) \subset {}^\sigma \mathbf{C}({}^*B)$ and in general ${}^\sigma(\mathbf{C}(B)) \neq ({}^\sigma \mathbf{C})({}^*B)$, ${}^\sigma(\mathbf{C}(F)) \neq ({}^\sigma \mathbf{C})(F)$.
- (iv) ${}^\sigma(\mathbf{C}(B)) = \bigcup \{ {}^\sigma(\mathbf{C}(F)) \mid F \in {}^\sigma(F(B)) \} = \bigcup \{ {}^\sigma(\mathbf{C}(F)) \mid F \in F({}^\sigma B) \} = \mathbf{C}(B) = \bigcup \{ \mathbf{C}(F) \mid F \in F(B) \}$, where $C \in \mathcal{C}_f$.

Throughout the remainder of this paper, we remove from $\mathcal{C}$ the one and only one inconsistent consequence operator U . Thus notationally we let $\mathcal{C}$ denote the set of all consequence operators defined on infinite $L \subset W$ with the exception of U . Two types of chains will be investigated. Let T be any chain in $\langle \mathcal{C}, \leq \rangle$ and T' be any chain with the additional property that for each $C \in T'$ there exists some $C' \in T'$ such that $C < C'$.

THEOREM 4.2 *There exists some $C_0 \in {}^*\mathbf{T}$ such that for each $C \in \mathbf{T}$, ${}^*\mathbf{C} \leq C_0$. There exists some $C'_0 \in {}^*\mathbf{T}'$ such that C'_0 is a purely subtle consequence operator and for each $C \in \mathbf{T}'$, ${}^*\mathbf{C} < C'_0$. Each member of ${}^*\mathbf{T}$ and ${}^*\mathbf{T}'$ are subtle consequence operators.*

PROOF. Let $R = \{(x, y) \mid x, y \in \mathbf{T} \text{ and } x \leq y\}$ and $R' = \{(x, y) \mid x, y \in \mathbf{T}' \text{ and } x < y\}$. In the usual manner, it follows that R and R' are concurrent on the set $\mathbf{T}$ and $\mathbf{T}'$ respectively. Thus there is some $C_0 \in {}^*\mathbf{T}$ and $C'_0 \in {}^*\mathbf{T}'$ such that for each $C \in \mathbf{T}$ and $C' \in \mathbf{T}'$, ${}^*\mathbf{C} \leq C_0$ and ${}^*\mathbf{C}' < C'_0$ since ${}^*\mathcal{M}$ is an enlargement. Note that the members of ${}^*\mathbf{T}$ and ${}^*\mathbf{T}'$ are defined on the set of all internal subsets of ${}^*\mathbf{L}$. However, if there is some similarly defined $D \in \mathcal{N}$ such that C_0 or $C'_0 = {}^\sigma D$, then since ${}^\sigma D$ is only defined for * -extensions of the (standard) members of $\mathcal{P}(\mathbf{L})$ and each $E \in {}^*\mathbf{T}$ or ${}^*\mathbf{T}'$ is defined on the internal subsets of ${}^*\mathbf{L}$ and there are internal subsets of ${}^*\mathbf{L}$ that are not * -extensions of standard sets we would have a contradiction. Of course, each member of ${}^*\mathbf{T}$ or ${}^*\mathbf{T}'$ is a subtle consequence operator. Hence each $E \in {}^*\mathbf{T}$ or ${}^*\mathbf{T}'$ is either equal to some ${}^*\mathbf{C}$, where $C \in \mathbf{T}$ or $C \in \mathbf{T}'$ or it is a purely subtle consequence operator. Now there does not exist a $D \in \mathcal{N}$ such that $C'_0 = {}^*\mathbf{D}$ since $C'_0 \in {}^*\mathbf{T}'$ and ${}^*\mathbf{C} \neq C'_0$ for each ${}^*\mathbf{C} \in {}^\sigma \mathbf{T}'$ would yield the contradiction that ${}^*\mathbf{D} \in {}^*\mathbf{T}' - {}^\sigma \mathbf{T}'$ but ${}^*\mathbf{D} \in {}^\sigma \mathcal{C}$. Hence C'_0 is a purely subtle consequence operator. This completes the proof.

Let $C \in \mathbf{T}'$. Since ${}^*\mathbf{C} < C'_0$, then C'_0 is “more powerful” than any $C \in \mathbf{T}'$ in the following sense. If $B \in \mathcal{P}(\mathbf{L})$, then for each $C \in \mathbf{T}'$ it follows that $\mathbf{C}(B) \subset {}^*(\mathbf{C}(B)) = {}^*\mathbf{C}({}^*\mathbf{B}) \subset C'_0({}^*\mathbf{B})$. Recall that, for $C \in \mathcal{C}$, a set $B \in \mathcal{P}(\mathbf{L})$ is called a *C-deductive system* if $C(B) = B$. From this point on, all results are restricted to chains in $\langle \mathcal{C}_f, \leq \rangle$.

THEOREM 4.3. *Let $C \in \mathbf{T} \cup \mathbf{T}'$ and $B \in \mathcal{P}(\mathbf{L})$. Then there exists a * -finite $F_0 \in {}^*(F(\mathbf{B}))$ such that $\mathbf{C}(B) \subset {}^*\mathbf{C}(F_0) \subset {}^*\mathbf{C}({}^*\mathbf{B}) = {}^*(\mathbf{C}(B))$ and ${}^*\mathbf{C}(F_0) \cap \mathbf{L} = \mathbf{C}(B) = {}^*\mathbf{C}(F_0) \cap \mathbf{C}(B)$.*

PROOF. Consider the binary relation $Q = \{(x, y) \mid x \in \mathbf{C}(B), y \in F(\mathbf{B}) \text{ and } x \in \mathbf{C}(y)\}$. By axiom (iii), the domain of Q is $\mathbf{C}(B)$. Let $(x_1, y_1), \dots, (x_n, y_n) \in Q$. By theorem 1 in [5, p. 64], (the monotone theorem) we have that $\mathbf{C}(y_1) \cup \dots \cup \mathbf{C}(y_n) \subset \mathbf{C}(y_1 \cup \dots \cup y_n)$. Since $F = y_1 \cup \dots \cup y_n \in F(\mathbf{B})$, then $(x_1, F), \dots, (x_n, F) \in Q$. Thus Q is concurrent on $\mathbf{C}(B)$. Hence there exists some $F_0 \in {}^*(F(\mathbf{B}))$ such that ${}^\sigma(\mathbf{C}(B)) = \mathbf{C}(B) \subset {}^*\mathbf{C}(F_0) \subset {}^*\mathbf{C}({}^*\mathbf{B}) = {}^*(\mathbf{C}(B))$. Since ${}^\sigma \mathbf{L} = \mathbf{L}$, then ${}^*\mathbf{C}(F_0) \cap \mathbf{L} = \mathbf{C}(B) = {}^*\mathbf{C}(F_0) \cap \mathbf{C}(B)$.

COROLLARY 4.3.1 *If $C \in \mathcal{C}_f$ and $B \in \mathcal{P}(\mathbf{L})$ is a C-deductive system, then there exists a * -finite $F_0 \subset {}^*\mathbf{B}$ such that ${}^*\mathbf{C}(F_0) \cap \mathbf{L} = B$.*

PROOF. Simply consider the one element chain $\mathbf{T} = \{C\}$.

COROLLARY 4.3.2. *Let $C \in \mathcal{C}_f$. Then there exists a * -finite $F_1 \subset {}^*\mathbf{L}$ such that for each C-deductive system $B \subset \mathbf{L}$, ${}^*\mathbf{C}(F_1) \cap B = B$.*

PROOF. Let $\mathbf{T} = \{C\}$ and the “B” in theorem 4.3 equal $\mathbf{L}$. The result now follows in a straightforward manner.

THEOREM 4.4. *Let $B \in \mathcal{P}(L)$.*

(i) *There exists a $*$ -finite $F_B \in {}^*(F(\mathbf{B}))$ and a subtle consequence operator $C_B \in {}^*\mathbf{T}$ such that for all $C \in \mathbf{T}$, ${}^\sigma(\mathbf{C}(\mathbf{B})) = \mathbf{C}(\mathbf{B}) \subset C_B(F_B)$.*

(ii) *There exists a $*$ -finite $F_B \in {}^*(F(\mathbf{B}))$ and a purely subtle consequence operator $C'_B \in {}^*\mathbf{T}'$ such that for all $C \in \mathbf{T}'$, ${}^\sigma(\mathbf{C}(\mathbf{B})) = \mathbf{C}(\mathbf{B}) \subset C'_B(F_B)$.*

PROOF. Consider the “binary” relation $Q = \{((x, z), (y, w)) \mid x \in \mathbf{T}, y \in \mathbf{T}, w \in F(\mathbf{B}), z \in x(w), z \in \mathcal{P}(\mathbf{L}), z \in x(w), \text{ and } x(w) \subset y(w)\}$. Let $\{((x_1, z_1), (y_1, w_1)), \dots, ((x_n, z_n), (y_n, w_n))\} \subset Q$. Notice that $F = w_1 \cup \dots \cup w_n \in F(\mathbf{B})$ and for the set $K = \{x_1, \dots, x_n\}$, let D be the largest member of K with respect to $\leq$. It follows that $z_i \in x_i(w_i) \subset x_i(F) \subset D(F)$ for each $i = 1, \dots, n$. Hence $\{((x_1, z_1), (D, F)), \dots, ((x_n, z_n), (D, F))\} \subset Q$ implies that Q is concurrent on its domain. Consequently, there exists some $(C_B, F_B) \in {}^*\mathbf{T} \times {}^*(F(\mathbf{B}))$ such that for each $(x, z) \in \text{domain of } Q$, $({}^*(x, z), (C_B, F_B)) \in {}^*Q$. Or, for each $(u, v) \in {}^\sigma(\text{domain of } Q)$, $((u, v), (C_B, F_B)) \in {}^*Q$. Let arbitrary $C \in \mathbf{T}$ and $b \in C(B)$. Then there exists some $F' \in F(\mathbf{B})$ such that $\mathbf{b} \in \mathbf{C}(F')$. Thus $({}^*\mathbf{C}, {}^*\mathbf{b}) \in {}^\sigma(\text{domain of } Q)$. Consequently, for each $C \in \mathbf{T}$ and $\mathbf{b} \in \mathbf{C}(\mathbf{B})$, $\mathbf{b} = {}^*\mathbf{b} \in ({}^*\mathbf{C})(F_B) \subset C_B(F_B)$. This all implies that for each $C \in \mathbf{T}$, ${}^\sigma(\mathbf{C}(\mathbf{B})) = \mathbf{C}(\mathbf{B}) \subset C_B(F_B)$.

(ii) Change the relation Q to Q' by requiring that $x \neq y$. Replace D in the proof of (i) above with D' is greater than and not equal to the largest member of K . Such a D' exists in $\mathbf{T}'$ from the definition of $\mathbf{T}'$. Continue the proof in the same manner in order to obtain C'_B and F'_B . The fact that C'_B is a purely subtle consequence operator follows in the same manner as in the proof of theorem 4.2.

COROLLARY 4.4.1 *There exists a [resp. purely] subtle consequence operator $C_L \in {}^*\mathbf{T}$ [resp. ${}^*\mathbf{T}'$] and a $*$ -finite $F_L \in {}^*(F(\mathbf{L}))$ such that for all $C \in \mathbf{T}$ [resp. $\mathbf{T}'$] and each $B \in \mathcal{P}(L)$, $\mathbf{B} \subset \mathbf{C}(\mathbf{B}) \subset C_L(F_L)$.*

PROOF. Simply let “ B ” in theorem 4.4 be equal to L . Then there exists a [resp. purely] subtle $C_L \in {}^*\mathbf{T}$ [resp. ${}^*\mathbf{T}'$] and $F_L \in {}^*(F(\mathbf{L}))$ such that for all $C \in \mathbf{T}$ [resp. $\mathbf{T}'$] $\mathbf{C}(\mathbf{L}) \subset C_L(F_L)$. If $B \in \mathcal{P}(L)$ and $C \in \mathbf{T}$ [resp. $\mathbf{T}'$], then $B \subset \mathbf{C}(\mathbf{B}) \subset \mathbf{C}(\mathbf{L})$. Thus for each $B \in \mathcal{P}(L)$ and $C \in \mathbf{T}$ [resp. $\mathbf{T}'$] $\mathbf{B} \subset \mathbf{C}(\mathbf{B}) \subset C_L(F_L)$ and the theorem is established.

The nonstandard results in this section have important applications to mathematical philosophy. We present two such applications. Let $\mathcal{F}$ be the symbolic alphabet for any formal language L with the usual assortment of primitive symbols [8, p. 59]. We note that it is possible to mimic the construction of L within $\mathcal{E}$ itself. If this is done, then it is not necessary to consider the intuitive map θ and we may restrict our attention entirely to the sets $\mathcal{E}$ and ${}^*\mathcal{E}$.

Let S denote the predicate consequence operator by the standard rules for predicate (proof-theory) deduction as they appear on pages 59 and 60 of reference [8]. Hence $A \in \mathcal{P}(L)$, $S(A) = \{x \mid x \in L \text{ and } A \vdash x\}$. It is not difficult to restrict the modus ponens rule of inference in such a manner that a denumerable set $\mathbf{T}' = \{C_n \mid n \in \mathbb{N}\}$ of consequence operators defined on $\mathcal{P}(L)$ is generated with the following properties.

- (i) For each $A \in \mathcal{P}(L)$, $S(A) = \bigcup \{C_n(A) \mid n \in \mathbb{N}\}$ and $C_n \neq S$ for any $n \in \mathbb{N}$.
- (ii) For each $C \in T'$ there is a C' such that $C < C'$ [4, p.57]. Let $A \in \mathcal{P}(L)$ be any S -deductive system. The $A = S(A) = \bigcup \{C_n(A) \mid n \in \mathbb{N}\}$ yields that A is a C_n -deductive system for each $n \in \mathbb{N}$. Thus S and C_n $n \in \mathbb{N}$ are consequence operators defined on $\mathcal{P}(A)$ as well as on $\mathcal{P}(L)$.

THEOREM 4.5. *Let L be a first-order language and $A \in \mathcal{P}(L)$. Then there exists a purely subtle $C_1 \in {}^*T'$ and a * -finite $F_1 \in {}^*(F(A))$ such that for each $B \in \mathcal{P}(A)$ and each $C \in T'$*

- (i) $\mathbf{C}(B) \subset C_1(F_1)$,
- (ii) $\mathbf{S}(B) \subset C_1(F_1) \subset {}^*\mathbf{S}(F_1) \subset {}^*(\mathbf{S}(A))$.
- (iii) ${}^*\mathbf{S}(F_1) \cap \mathbf{L} = \mathbf{S}(A) = C_1(F_1) \cap \mathbf{L}$.

PROOF. The same proof as for corollary 4.4.1 yields that there is some purely subtle $C_1 \in {}^*T'$ and $F_1 \in {}^*(F(A))$ such that for each $B \in \mathcal{P}(A)$ and each $C \in T'$, $\mathbf{C}(B) \subset C_1(F_1)$ and (i) follows. From (i), it follows that $\bigcup \{\sigma(\mathbf{C}(B)) \mid C \in T'\} = \bigcup \{\mathbf{C}(B) \mid C \in T'\} = \mathbf{S}(B) = \sigma(\mathbf{S}(B)) \subset C_1(F_1)$ and the first part of (ii) holds. By * -transfer $C_1 < {}^*\mathbf{S}$ and C_1 and ${}^*\mathbf{S}$ are defined on internal subsets of ${}^*\mathbf{A}$. Thus $C_1(F_1) \subset {}^*\mathbf{S}(F_1) \subset {}^*\mathbf{S}({}^*\mathbf{A}) = {}^*(\mathbf{S}(A))$ by the * -monotone property. This completes (ii). Since $\mathbf{S}(A) \subset C_1(F_1) \subset {}^*\mathbf{S}(F_1) \subset {}^*(\mathbf{S}(A))$ from (ii), then (iii) follows and the theorem is proved.

REMARK 4.6. Of course, it is well known that there exists some $F \in {}^*(F(A))$ such that $\mathbf{S}(A) \supset \mathbf{A} \subset F \subset {}^*\mathbf{A}$ and * -transfer of axiom (i) yields that ${}^*\mathbf{S}(F) \subset {}^*\mathbf{S}({}^*\mathbf{A}) = {}^*(\mathbf{S}(A))$. However, F_1 of theorem 4.5 is of a special nature in that the purely subtle C_1 applied to F_1 yields the indicated properties. Also theorem 4.5 holds for many other infinite languages and deductive processes.

Let L be a language and let M be a structure in which L can be interpreted in the usual manner. A consequence operator C is *sound* for M if whenever $A \in \mathcal{P}(L)$ has the property that $M \models A$, then $M \models C(A)$. As usual, $T(M) = \{x \mid x \in L \text{ and } M \models x\}$. Obviously, if C is sound for M , then $T(M)$ is a C -deductive system.

Corollary 4.3.1 implies that there exists * -finite $F_0 \subset {}^*(T(M))$ such that ${}^*\mathbf{C}(F_0) \cap \mathbf{L} = T(M)$. Notice that the fact that F_0 is * -finite implies that F_0 is * -recursive. Moreover, trivially, F_0 is a * -axiom system for ${}^*\mathbf{C}(F_0)$, and we do not lack knowledge about the behavior of F_0 since any formal property about $\mathbf{C}$ or recursive sets, among others, must hold for ${}^*\mathbf{C}$ or F_0 when property interpreted. If L is a first-order language, then S is sound for first-order structures. Theorem 4.5 not only yields a * -finite F_1 but a purely subtle consequence operator C_1 such that, trivially, F_1 is a * -axiom for $C_1(F_1)$ and for ${}^*\mathbf{S}(F_1)$. In this case, we have that ${}^*\mathbf{S}(F_1) \cap \mathbf{L} = T(M) = C_1(F_1) \cap \mathbf{L}$. By the use of internal and external objects, the nonstandard logics $\{ {}^*\mathbf{C}, {}^*\mathbf{L} \}$, $\{ C_1, {}^*\mathbf{L} \}$ and $\{ {}^*\mathbf{S}, {}^*\mathbf{L} \}$ technically by-pass a portion of Gödel's first incompleteness theorem.

By definition $b \in S(B)$, $B \in \mathcal{P}(L)$ iff there is a finite length “proof” of b from the premises B . It follows, that for each $b \in {}^*(T(M))$ there exists a * -finite length proof of b from a * -finite set of premises F_1 . If we let ${}^*\mathcal{M}$ be an enlargement with

the $\aleph_1$ -isomorphism property, among others, then each *-finite length proof is either externally finite or externally infinite, and all externally infinite proof lengths are of the same external cardinality [3].

References

1. W. Dziobiak: The lattice of strengthenings of a strongly finite consequence operator, *Studia Logica*, **40**(2) (1981), 177-193.
2. J. R. Geiser: Nonstandard logics, *J. Symbolic Logic*, **33**(1968), 236-250.
3. C. W. Henson: The isomorphism property in nonstandard analysis and its use in the theory of Banach spaces, *J. Symbolic Logic*, **39**(1974), 717-731.
4. R. A. Herrmann: The mathematics for mathematical philosophy, Monograph #130, Institute for Mathematical Philosophy Press, Annapolis, 1983. (Incorporated into "The Theory of Ultralogics," <http://xxx.lanl.gov/abs/math.GM/9903081> and 9903082.)
5. R. A. Herrmann: Mathematical Philosophy, *Abstracts A. M. S.* **2**(6)(1981), 527.
6. J. Los and R. Suszko: Remarks on sentential logics, *Indag. Math.*, **20**(1958), 177-183.
7. A. A. Markov: Theory of algorithms, *Amer. Math. Soc. Translations, Ser. 2*, **15**(1960), 1-14.
8. E. Mendelson: Introduction to mathematical logic, D. Van Nostrand, New York, 1979.
9. A. Robinson: On languages which are based on non-standard arithmetic, *Nagoya Math. J.*, **22**(1963), 83-118.
10. A. Tarski: Logic, semantics and metamathematics, Oxford University Press, Oxford 1956.
11. R. Wójcicki: Some remarks on the consequence operation in sentential logics, *Fund. Math.* **68**(1970), 269-279.

Robert A. Herrmann
 Mathematics Department,
 U. S. Naval Academy,
 ANNAPOLIS, MD 21402
 U. S.

The material that begins on the next page (23) is taken from my more comprehensive book "The Theory of Ultralogics" (Ultralogics and More) with a minor addition or two. In particular, the additional section 11.2 does not appear in that article. On the other hand, the entire mathematical foundations and additional material that leads to the G and D-world models can be found in "The Theory of Ultralogics."

7.1 Introduction.

Consider the real line. If you believe that time is the ordinary continuum, then the entire real line can be your time line. Otherwise, you may consider only a subset of the real line as a time line. In the original version of this section, the time concept for the GGU-model was presented in an unnecessarily complex form. As shown in [3], one can assume an absolute substratum time within the NSP-world. It is the infinitesimal light-clock time measures that may be altered by physical processes. In my view, the theory of quantum electrodynamics would not exist without such a NSP-world time concept.

Consider a small interval $[a, b]$, $a < b$ as our basic time interval where as the real numbers increase the time is intuitively considered to be increasing. In the following approach, one may apply the concept of the persistence of mental version relative to descriptions for the behavior of a Natural-system at a moment of time within this interval. An exceptionally small subinterval can be chosen within $[a, b]$ as a maximum subinterval length $= M$. “Time” and the size of a “time” interval as they are used in this and the following sections refer to an intuitive concept used to aid in comprehending the notation of an event sequence - an event ordering concept. First, let $a = t_0$. Then choose t_1 such that $a < t_1 < b$. There is a partition $t_1, \dots, t_m$ of $[a, b]$ such that $t_0 < t_1 < \dots < t_m < b$ and $t_{j+1} - t_j \leq M$. The final subinterval $[t_m, b]$ is now separated, by induction, say by taking midpoints, into an increasing sequence of times $\{t_q\}$ such that $t_m < t_q < b$ for each q and $\lim_{q \rightarrow \infty} t_q = b$.

Assume the prototype $[a, b]$ with the time subintervals as defined above. Let $[t_j, t_{j+1})$ be any of the time subintervals in $[a, b]$. For each such subinterval, let W_i denote the readable sentence

This||frozen||segment||gives||a||description||for||the||
time||interval||that||has||as||its||leftmost||endpoint||the
||time|| $\lceil t_i \rceil$ ||that||corresponds||to||the||natural||number|| i .

Let $T_i = \{xW_i \mid x \in \mathcal{W}\}$. The set T_i is called a *totality* and each member of any such T_i is called a *frozen segment*. Notice that since the empty word is not a member of $\mathcal{W}$, then the cardinality of each member of T_i is greater than that of W_i . Each T_i is a (Dedekind) denumerable set, and if $i \neq j$, then $T_i \cap T_j = \emptyset$.

W_i is only an identifier and may be altered. “Time,” either its measure or otherwise, is not the actual underlying interpretation for these intervals. Time refers to an external event ordering concept, an intuitive *event sequence*. For most purposes, simply call these intervals “event intervals.” [These event intervals can correspond to the universal event numbers concept.] In the above descriptions for W_i , simply replace “time||interval” with “event||interval” and replace the second instance of the word “time” with the word “event.” If this event sequence interpretation is made, then other compatible interpretations would be necessary when applying some of the following results.

I point out two minor aspects of the above constructions. First, within certain descriptions there are often “symbols” used for real, complex, natural numbers etc. These objects also exist as abstract objects within the structure $\mathcal{M}$. No inconsistent interpretations should occur when these objects are specifically modeled within $\mathcal{M}$ since to my knowledge all of the usual mathematical objects used within physical analysis are disjoint from $\mathcal{E}$ as well as disjoint from any finite Cartesian product of $\mathcal{E}$ with itself. If for future research within physical applications finite partial sequences of natural numbers and the finite equivalence classes that appear in $\mathcal{E}$ are needed and are combined into one model for different purposes than the study of descriptions, then certain modifications would need to

be made so that interpretations would remain consistent. Secondly, I have tried whenever intuitive strings are used or sets of such strings are defined to use Roman letter notation for such objects. This only applies for the intuitive model. Also W_i is only an identifier and may be altered.

7.2 Developmental Paradigms

It is clear that if one considers a time interval of the type $(-\infty, +\infty)$, $(-\infty, b)$ or $[a, +\infty)$, then each of these may be considered as the union of a denumerable collection of time intervals of the type $[a, b)$ with common endpoint names displayed. Further, although $[a, b)$ is to be considered as subdivided into denumerably many subintervals, it is not necessary that each of the time intervals $[t_j, t_{j+1}) \subset [a, b)$ be accorded a corresponding description for the appearance of a specific Natural-system that is distinct from all others that occur throughout the time subinterval. Repeated descriptions only containing a different last natural number i in the next to last position will suffice. Each basic developmental paradigm will be restricted, at present, to such a time interval $[a, b)$.

Where human perception and descriptive ability is concerned, the least controversial approach would be to consider only finitely many descriptive choices as appropriate. A finite set is recursive and such a choice, since the result is such a set, would be considered to be the simplest type of algorithm. You “simply” check to see if an expression is a member of such a finite set. If we limited ourselves to finitely many human choices for Natural system descriptions from the set of all totalities and did not allow a denumerable or a continuum set to be chosen, then the next result establishes that within the Nonstandard Physical world (i.e. NSP-world) such a finite-type of choice can be applied and a continuum of descriptions obtained.

The following theorem is not insignificant even if we are willing to accept a denumerable set of distinct descriptions — descriptions that are not only distinct in the next to the last symbol, but are also distinctly different in other aspects as well. For, if this is the case, the results of Theorem 7.2.1 still apply. The same finite-type of process in the NSP-world yields such a denumerable set as well.

The term “NSP-world” will signify a certain second type of interpretation for nonstandard entities. In particular, the subtle logics, unreadable sentences, etc. This interpretation will be developed throughout the remainder of this book. One important aspect of how descriptions are to be interpreted is that a description correlates directly to an assumed or observed real Natural phenomenon, and conversely. In these investigations, the phenomenon is called an *event*.

In order to simplify matters a bit, the following notation is employed. Let $\mathcal{T} = \{T_i \mid i \in \mathbb{N}\}$. Let $F(\mathcal{T})$ be the set of all **nonempty** and *finite* subsets of $\mathcal{T}$. This symbol has been used previously to include the empty set, this set is now excluded. Now let $A \in F(\mathcal{T})$. Then there exists a finite choice set s such that $x \in s$ iff there exists a unique $T_i \in A$ and $x \in T_i$. Now let the set $\mathcal{C}$ denote the set of all such finite choice sets. As to interpreting these results within the NSP-world, the following is essential. Within nonstandard analysis the term “hyper” is often used for the result of the $*$ map. For example, you have ${}^*\mathbb{R}$ as the hyperreals since $\mathbb{R}$ is termed the real numbers. For certain, but not all concepts, the term “hyper” or the corresponding $*$ notation will be universally replaced by the term “ultra.” Thus, certain purely subtle words or $*$ -words become “ultrawords” within the developmental paradigm interpretation. [Note: such a word was previously called a superword.] Of course, for other scientific or philosophical systems, such abstract mathematical objects can be reinterpreted by an appropriate technical term taken from those disciplines.

As usual, we are working within any enlargement and all of the above intuitive objects are embedded into the G-structure. Recall, that to simplify expressions, we often suppress within our

first-order statements a specific superstructure element that bounds a specific quantifier. [Note 2 MAY 1998: The material between the [[and the]] has been altered from the original that appeared in the 1993 revision.] [[Although theorem 7.2.1 may not be insignificant, it is also not necessary for the other portions of this research. The general axiom of choice can be applied to generate formally developmental paradigms.

The $\mathcal{S}$ in Theorem 7.2.1 determines a special intrinsic ultranatural selection process (i.e. **IUN-selection process**). This process is obtained by application of hyperfinite choice.

Theorem 7.2.1 Let $\emptyset \neq \gamma \subset \mathbb{N}$ and $\tilde{\mathcal{T}} = \{\mathbf{T}_i \mid i \in \gamma\}$. There exists a set of sets $\mathcal{S}$ determined by hyperfinite set Q and hyper finite choice defined on Q such that:

(i) $s' \in \mathcal{S}$ iff for each $\mathbf{T} \in \tilde{\mathcal{T}}$ there is one and only one $[g] \in {}^*\mathbf{T}$ such that $[g] \in s'$, and if $x \in s'$, then there is some $\mathbf{T} \in \tilde{\mathcal{T}}$ and some $[g] \in {}^*\mathbf{T}$ such that $x = [g]$. (If ${}^*[g] \in {}^\sigma\mathbf{T}$, then $[g] = [f] \in \mathbf{T}$.)

Proof. (i) Let $A \in F(\tilde{\mathcal{T}})$. Then from the definition of $\tilde{\mathcal{T}}$, there exists some $n \in \mathbb{N}$ such that $A = \{\mathbf{T}_{j_i} \mid i = 0, \dots, n \wedge j_i \in \mathbb{N}\}$. From the definition of $\mathbf{T}_k$, each $\mathbf{T}_k$ is denumerable. Notice that any $[f] \in \mathbf{T}_k$ is associated with a unique member of A_1 . Simply consider the unique $f_0 \in [f]$. The unique member of A_1 is by definition $f_0(0)$. Thus each member of $\mathbf{T}_k$ can be specifically identified. Hence, for each $\mathbf{T}_i$ there is a denumerable $M_i \subset \mathbb{N}$ and a bijection $h_i: M_i \rightarrow \mathbf{T}_i$ such that $a_i \in \mathbf{T}_i$ iff there is a $k_i \in M_i$ and $h_i(k_i) = a_i$. Consequently, for each $i = 0, \dots, n$ and $a_{j_i} \in \mathbf{T}_{j_i}$, we have that $h_{j_i}(k_{j_i}) = a_{j_i}$, and conversely for each $i = 0, \dots, n$ and $k_{j_i} \in M_{j_i}$, $h_{j_i}(k_{j_i}) \in \mathbf{T}_{j_i}$. Obviously, $\{h_{j_i}(k_{j_i}) \mid i = 0, \dots, n\}$ is a finite choice set. All of the above may be translated into the following sentence that holds in $\mathcal{M}$. (Note: Choice sets are usually considered as the range of choice functions. Further, “bounded formula simplification” has been used.)

$$\begin{aligned} \forall y(y \in F(\tilde{\mathcal{T}}) \rightarrow \exists s((s \in \mathcal{P}(\mathcal{E})) \wedge \forall x((x \in y) \rightarrow \exists z((z \in x) \wedge (z \in s) \wedge \\ \forall w(w \in \mathcal{E} \rightarrow ((w \in s) \wedge (w \in x) \leftrightarrow (w = z)))))) \wedge \\ \forall u(u \in \mathcal{E} \rightarrow ((u \in s) \leftrightarrow \exists x_1((x_1 \in y) \wedge (u \in x_1)))))) \end{aligned} \quad (7.2.1)$$

For each $A \in F(\tilde{\mathcal{T}})$, let S_A be the set of all such choice sets generated by the predicate that follows the first $\rightarrow$ formed from (7.2.1) by deleting the $\exists s$ and letting $y = A$. Of course, this set exists within our set theory. Now let $\mathcal{C} = \{S_A \mid A \in F(\tilde{\mathcal{T}})\}$.

Consider ${}^*\mathcal{C}$ and ${}^*(S_A)$. Then $s \in {}^*(S_A)$ iff s satisfies (7.2.1) as interpreted in ${}^*\mathcal{M}$. Since we are working in an enlargement, there exists an internal $Q \in {}^*(F(\tilde{\mathcal{T}}))$ such that ${}^\sigma\tilde{\mathcal{T}} \subset Q \subset {}^*\tilde{\mathcal{T}}$. Recall that ${}^\sigma\tilde{\mathcal{T}} = \{{}^*\mathbf{T} \mid \mathbf{T} \in \tilde{\mathcal{T}}\}$. Also ${}^\sigma\mathbf{T} \subset {}^*\mathbf{T}$ for each $\mathbf{T} \in \tilde{\mathcal{T}}$. From the definition of ${}^*\mathcal{C}$, there is an internal set S_Q and $s \in S_Q$ iff s satisfies the internal defining predicate for members of S_Q and this set is the set of all such s . ($\Rightarrow$) Consequently, since for each $\mathbf{T} \in \tilde{\mathcal{T}}$, ${}^*\mathbf{T} \in Q$, then the generally external $s' = \{s \cap {}^*\mathbf{T} \mid \mathbf{T} \in \tilde{\mathcal{T}}\}$ satisfies the $\Rightarrow$ for (i). Note, however, that for ${}^*\mathbf{T}$, $\mathbf{T} \in \tilde{\mathcal{T}}$, it is possible that $s \cap {}^*\mathbf{T} = \{{}^*[f]\}$ and ${}^*[f] \in {}^\sigma\mathbf{T}$. In this case, by the finiteness of $[f]$ it follows that $[f] = {}^*[f]$ implies that $s \cap {}^*\mathbf{T} = \{[f]\}$. Now let $\mathcal{S} = \{s' \mid s \in S_Q\}$. In general, $\mathcal{S}$ is an external object.

($\Leftarrow$) Consider the internal set S_Q . Let s' be the set as defined by the right-hand side of (i). For each internal $x \in s'$ and applying, if necessary, the * -axiom of choice for * -finite sets, we have the internal set $A_x = \{y \mid (y \in S_Q) \wedge (x \in y)\}$ is nonempty. The set $\{A_x \mid x \in s'\}$ has the finite intersection property. For, let nonempty internal $B = \{x_1, \dots, x_n\}$. Then the set $A_B = \{y \mid (y \in S_Q) \wedge (x_1 \in y) \cdots \wedge (x_n \in y)\}$ is internal and nonempty by the * -axiom of choice for * -finite sets. Since we are in an enlargement and s' is countable, then $D = \bigcap \{A_x \mid x \in s'\} \neq \emptyset$. Now take any $s \in D$. Then $s \in S_Q$ and from the definition of $\mathcal{S}$, $s' \in \mathcal{S}$. This completes the proof. ■

[Note: Theorem 7.2.1 may be used to model physical developmental paradigms associated with event sequences.]

Although it is not necessary, for this particular investigation, the set $\mathcal{S}$ may be considered a *set of all developmental paradigms*. Apparently, $\mathcal{S}$ contains every possible developmental paradigm for all possible frozen segments and $\mathcal{S}$ contains paradigms for any $*$ -totality $*\mathbf{T}$. There are $*$ -frozen segments contained in various s' that can be assumed to be unreadable sentences since $\sigma\mathbf{T} \neq *\mathbf{T}$.]

Let $A \in F(\tilde{\mathcal{T}})$ and $M(A)$ be a subset of S_A for which there exists a written set of rules that selects some specific member of S_A . Obviously, this may be modeled by means of functional relations. First, $M(A) \subset S_A$ and it follows, from the difference in cardinalities, that there are infinitely many members of $*(S_A)$ for which there does not exist a readable rule that will select such members. However, this does not preclude the possibility that there is a set of purely unreadable sentences that do determine a specific member of $*S_A - \sigma M(A)$. This might come about in the following manner. Suppose that H is an infinite set of formal sentences that is interpreted to be a set of rules for the selection of distinct members of $M(A)$. Suppose we have a bijection $h: M(A) \rightarrow \mathbf{H}$ that represents this selection process. Let $*\mathcal{M}$ be at least a polysaturated enlargement of $\mathcal{M}$, and consider $\sigma f: \sigma(M(A)) \rightarrow \sigma\mathbf{H}$. The map σf is also a bijection and $\sigma f: \sigma(M(A)) \rightarrow *\mathbf{H}$. Since $|\sigma(M(A))| < |\mathcal{M}|$, it is well-known that there exists an internal map $h: A' \rightarrow *\mathbf{H}$ such that $h \mid \sigma(M(A)) = \sigma f$, and A' , $h[A']$ are internal. Further, for internal $A' \cap *(S_A) = B$, $\sigma(M(A)) \subset B$. However, $\sigma(M(A))$ is external. This yields that h is defined on B and $B \cap (*S_A - \sigma(M(A))) \neq \emptyset$. Also, $\sigma\mathbf{H} \subset h[B] \subset *\mathbf{H}$ implies, since $h[B]$ is internal, that $\sigma H \neq h[B]$. Consequently, in this case, $h[B]$ may be interpreted as a set of $*$ -rules that determine the selection of members of B . That is to say that there is some $[g] \in h[B] - \sigma H$ and a $[k] \in *S_A - \sigma(M(A))$ such that $([k], [g]) \in h$. As it will be shown in the next section, the set H can be so constructed that if $[g] \in h[B] - \sigma H$, then $[g]$ is unreadable.

7.3 Ultrawords

Ordinary propositional logic is not compatible with deductive quantum logic, intuitionistic logic, among others. In this section, a subsystem of propositional logic is investigated which rectifies this incompatibility. I remark that when a standard propositional language L or an informal language P isomorphic to L is considered, it will always be the case that the L or P is minimal relative to its applications. This signifies that if L or P is employed in our investigation for a developmental paradigm, then L or P is constructed only from those distinct propositional atoms that correspond to distinct members of d , etc. The same minimizing process is always assumed for the following constructions.

Let B be a formal or, informal nonempty set of propositions. Construct the language P_0 in the usual manner from B (with superfluous parentheses removed) so that P_0 forms the smallest set of formulas that contains B and such that P_0 is closed under the two binary operators $\wedge$ and $\rightarrow$ as they are formally or informally expressed. Of course, this language may be constructed inductively or by letting P_0 be the intersection of all collections of such formula closed under $\wedge$ and $\rightarrow$.

We now define the deductive system S . Assume substitutivity, parenthesis reduction and the like. Let $d = \{F_i \mid i \in \mathbb{N}\} = B$ be a development paradigm, where each F_i is a readable frozen segment and describes the behavior of a Natural-system over a time subinterval. Let the set of axioms be the schemata

$$(\mathcal{A} \wedge \mathcal{B}) \rightarrow \mathcal{A}, \mathcal{A} \in \mathcal{B} \quad (1)$$

$$(\mathcal{A} \wedge \mathcal{B}) \rightarrow \mathcal{B} \quad (2)$$

$$\mathcal{A} \wedge (\mathcal{B} \wedge \mathcal{C}) \rightarrow (\mathcal{A} \wedge \mathcal{B}) \wedge \mathcal{C}, \quad (3)$$

$$(\mathcal{A} \wedge \mathcal{B}) \wedge \mathcal{C} \rightarrow \mathcal{A} \wedge (\mathcal{B} \wedge \mathcal{C}). \quad (4)$$

If P_0 is considered as informal, which appears to be necessary for some applications, where the parentheses are replaced by the concept of symbol strings being to the “left” or “right” of other symbol strings and the concept of strengths of connectives is used (i.e. $A \wedge B \rightarrow C$ means $((A \wedge B) \rightarrow C)$), then axioms 3 – 4 and the parentheses in (1) and (2) may be omitted. The one rule of inference is Modus Ponens (MP). Proofs or demonstrations from hypotheses Γ contain finitely many steps, hypotheses may be inserted as steps and the last step in the proof is either a theorem if $\Gamma = \emptyset$ or if $\Gamma \neq \emptyset$, then the last step is a consequence of (a deduction from) Γ . Notice that repeated application of (4) along with (MP) will allow all left parentheses to be shifted to the right with the exception of the (suppressed) outermost left one. Thus this leads to the concept of left to right ordering of a formula. This allows for the suppression of such parentheses. In all the following, this suppression will be done and replaced with formula left to right ordering.

For each $\Gamma \subset P_0$, let $S(\Gamma)$ denote the set of all formal theorems and consequences obtained from the above defined system S . Since hypotheses may be inserted, for each $\Gamma \subset P_0$, $\Gamma \subset S(\Gamma) \subset P_0$. This implies that $S(\Gamma) \subset S(S(\Gamma))$. So, let $A \in S(S(\Gamma))$. The general concept of combining together finitely many steps from various proofs to yield another formal proof leads to the result that $A \in S(\Gamma)$. Therefore, $S(\Gamma) = S(S(\Gamma))$. Finally, the finite step requirement also yields the result that if $A \in S(\Gamma)$, then there exists a finite $F \subset \Gamma$ such that $A \in S(F)$. Consequently, S is a finitary consequence operator and observe that if C is the propositional consequence operator, then $S(\Gamma) \subsetneq C(\Gamma)$. Of course, we may now apply the nonstandard theory of consequence operators to S .

It is well-known that the axiom schemata chosen for S are theorems in intuitionistic logic. Now consider quantum logic with the Mittelstaedt conditional $i_1(A, B) = A^\perp \vee (A \wedge B)$. [1] Notice that $i_1(A \wedge B, B) = (A \wedge B)^\perp \vee ((A \wedge B) \wedge B) = (A \wedge B)^\perp \vee (A \wedge B) = I$ (the upper unit.) Then $i_1((A \wedge B), A) = (A \wedge B)^\perp \vee ((A \wedge B) \wedge A) = (A \wedge B)^\perp \vee (A \wedge B) = I$; $i_1((A \wedge B) \wedge C, A \wedge (B \wedge C)) = ((A \wedge B) \wedge C)^\perp \vee (A \wedge (B \wedge C)) = I = i_1(A \wedge (B \wedge C), (A \wedge B) \wedge C)$. Thus with respect to the interpretation of $\mathcal{A} \rightarrow \mathcal{B}$ as conditional i_1 the axiom schemata for the system S are theorems and the system S is compatible with deductive quantum logic under the Mittelstaedt conditional.

In what follows, assume $\wedge$ is interpreted as $|||and|||$. Recall that $d = \{F_i \mid i \in \mathbb{N}\} = B$ is a development paradigm, where each F_i is a readable frozen segment and describes the behavior of a Natural-system over a time subinterval. Although it is sufficient in most cases to consider a formal language, it is more convenient to employ an isomorphic informal language. This is especially necessary if one wishes to more closely analyze certain special subtle objects. Let $M_0 = d$. Define $M_1 = \{F_0|||and|||F_1\}$. Assume that M_n is defined. Define $M_{n+1} = \{x|||and|||F_{n+1} \mid x \in M_n\}$. From the fact that d is a developmental paradigm, where the last two symbols in each member of d is the time indicator “i.”, it follows that no member of d is a member of M_n for $n > 0$. Now let $M = \bigcup\{M_n \mid n \in \mathbb{N}\}$. Intuitively, $|||and|||$ behaves as a conjunction and each F_i as an atom within our language. Notice the important formal demonstration fact that for an hypothesis consisting of any member of M_n , $n > 0$, repeated applications of (1), (MP), (2), (MP) will lead to the members of d appearing in the proper time ordering at increasing (formal) demonstration step numbers.

The next theorem shows the existence of the most basic ultraword w . The ${}^*\mathbf{S}$ is the basic compatible ultralogic. This ultraword and the corresponding ultralogic generate a selected Natural event sequence along with numerous ultranatural events.

Theorem 7.3.1 For $\mathbf{d} = \{\mathbf{F}_i \mid i \in \mathbb{N}\}$, there exists an ultraword $w \in {}^*\mathbf{M} - {}^*\mathbf{d}$ such that $\mathbf{F}_i \in {}^*\mathbf{S}(\{w\})$ (i.e. $w \vdash_S \mathbf{F}_i$) for each $i \in \mathbb{N}$.

Proof. Consider the binary relation $G = \{(x, y) \mid (x \in \mathbf{d}) \wedge (y \in \mathbf{M} - \mathbf{d}) \wedge (x \in \mathbf{S}(\{y\}))\}$. Suppose that $\{(x_1, y_1), \dots, (x_n, y_n)\} \subset G$. For each $i = 1, \dots, n$ there is a unique $k_i \in \mathbb{N}$ such that $x_i = \mathbf{F}_{k_i}$. Let $m = \max\{k_i \mid (x_i = \mathbf{F}_{k_i}) \wedge (i = 1, \dots, n)\}$. Let $b \in \mathbf{M}_{m+1}$. It follows immediately that $x_i \in \mathbf{S}(\{b\})$ for each $i = 1, \dots, n$ and, from the construction of $\mathbf{d}$, $b \notin \mathbf{d}$. Thus $\{(x_1, b), \dots, (x_n, b)\} \subset G$. Consequently, G is a concurrent relation. Hence, there exists some $w \in {}^*\mathbf{M} - {}^*\mathbf{d}$ such that ${}^\sigma\mathbf{F}_i = \mathbf{F}_i \in {}^*\mathbf{S}(\{w\})$ for each $i \in \mathbb{N}$. This completes the proof. [See note 3.]

Observe that w in Theorem 7.3.1 has all of the formally expressible properties of a readable word. For example, w has a hyperfinite length, among other properties. However, since $\mathbf{d}$ is a denumerable set, each ultraword has a very special property.

Recall that for each $[g] \in \mathcal{E}$ there exists a unique $m \in \mathbb{N}$ and $f' \in T^m$ such that $[f'] = [g]$ and for each k such that $m < k \in \mathbb{N}$, there does not exist $g' \in T^k$ such that $[g'] = [g]$. The function $f' \in T^m$ determines all of the alphabet symbols, the symbol used for the blank space, and the like, and determines there position within the intuitive word being represented by $[g]$. Also for each j such that $0 \leq j \leq m$, $f'(j) = i(a) \in i[\mathcal{W}] = T \subset A_1$, where $i(a)$ is the “encoding” in A_1 of the symbol “a”. For each $m \in \mathbb{N}$, let $P_m = \{f \mid (f \in T^m) \wedge (\exists z((z \in \mathcal{E}) \wedge (f \in z) \wedge \forall x((x \in \mathbb{N}) \wedge (x > m) \rightarrow \neg \exists y((y \in T^x) \wedge (y \in z))))))\}$. An element $n \in {}^*A_1$ is a *subtle alphabet symbol* if there exists $m \in \mathbb{N}$ and $f \in {}^*(P_m)$, or if $m \in \mathbb{N}_\infty$ and $f \in P_m$, and some $j \in {}^*\mathbb{N}$ such that $f(j) = n$. A symbol is a *pure subtle alphabet symbol* if $f(j) = n \notin i[\mathcal{W}]$. Subtle alphabet symbols can be characterized in ${}^*\mathcal{E}$ for they are singleton objects. A $[g] \in {}^*\mathcal{E}$ represents a subtle alphabet symbol iff there exists some $f \in ({}^*T)^0$ such that $[f] = [g] = [(0, f(0))]$, $f = \{(0, f(0))\}$.

Theorem 7.3.2 Let $\mathbf{d} = \{\mathbf{F}_i \mid i \in \mathbb{N}\}$ be a denumerable developmental paradigm. For an ultraword $w = [g] \in {}^*\mathcal{E}$ that exists by Theorem 7.3.1, there exists $\delta \in \mathbb{N}_\infty$ and infinitely many $\nu \in \mathbb{N}_\infty$ such that $[f] = [g]$, $f \in P_\delta$ and each $f(\nu) \in {}^*(i[\mathbf{P}_0]) - (i[\mathbf{P}_0])$ is a pure subtle alphabet symbol.

Proof. Since $\mathbf{d}$ is denumerable, consider a bijection $h: \mathbb{N} \rightarrow \mathbf{d}$ such that $h(n) = \mathbf{F}_n = [(0, q_n)]$, $q_n \in i[\mathbf{d}] \subset i[\mathbf{P}_0]$. From the definition of $\mathbf{M} - \mathbf{d}$, if $[g] \in \mathbf{M} - \mathbf{d}$, there exists a unique m , $n \in \mathbb{N}$ ($n \geq 1$) and $f' \in (i[\mathbf{P}_0])^m$ such that $h[[0, n]] \subset \mathbf{S}(\{[g]\})$ and **(0)** $[f'] = [g]$, and **(1)** $m = 2n$, **(2)** for each even $2k \leq m$, ($k \geq 0$), $f'(2k) = q_k \in i[\mathbf{P}_0] \subset A_1$, $[(0, q_k)] \in \mathbf{d}$. All such q_k are distinct and $[(0, q_k)] \in \mathbf{S}(\{f'\})$. For each odd $2k+1 \leq m$, $f'(2k+1) = i(\| \text{and} \|)$. **(3)** For each $h(k) \in h[[0, n]]$ there exist distinct even $2k \leq m$ such that $f'(2k) = q_k \in i[\mathbf{P}_0] \subset A_1$ and conversely, and $h(k) = [(0, q_k)] \in \mathbf{d}$ all such q_k being distinct. Also note that for each $i \in \mathbb{N}$, $h(i) \in \mathbf{S}(\{[g]\})$ iff $h(i) \in h[[0, n]]$. **(4)** There exists a unique $r \in \mathbb{N}$ such that $r > m$ and an $f'' \in P_r$ such that $[f''] = [g]$. **(5)** For each of the $n+1$ distinct k 's that exist from the first part of **(2)**, there exists at least $n+1$ distinct $r_k \in \mathbb{N}$ such that $0 \leq r_n \leq r$ and $n+1$ distinct $f''(r_k) = i(b_k)$, where b_k is a symbol representing the natural number that appears as the next to the last symbol in a member of $\mathbf{d}$.

Let $w = [g] \in {}^*\mathbf{M} - {}^*\mathbf{d}$ be the ultraword that exists by Theorem 7.3.1. Then there exists some $\delta \in {}^*\mathbb{N}$ and a unique $\nu \in {}^*\mathbb{N}$ as well as $f' \in ({}^*i[\mathbf{P}_0])^\delta$ such that ${}^*h[[0, \nu]] \subset {}^*\mathbf{S}(\{f'\}) = {}^*\mathbf{S}(\{[g]\})$

with the $*$ -transfer of properties (1) — (5). From Theorem 7.3.1, $\sigma(h[\mathbb{N}]) \subset {}^*\mathbf{S}(\{[f']\})$ yields, by application, of (3) that $h[\mathbb{N}] = \sigma(h[\mathbb{N}]) \subset {}^*h[[0, \nu]]$. This implies that ${}^*h[[0, \nu]]$ is infinite and internal. From (1), we obtain that $\delta \in \mathbb{N}_\infty$. From (4), there exists a unique $\rho > \delta$, hence $\rho \in \mathbb{N}_\infty$, and a unique $f \in P_\delta$ such that $[f] = [g] = [f']$. From (5), there exist at least ν $*$ -distinct, and hence distinct, $\rho_k \in {}^*\mathbb{N}$ such that the ν distinct $f(\rho_k) \in {}^*(i[P_0])$. Since ${}^*\mathcal{M}$ is an ultrapower or ultralimit enlargement based upon $\mathbb{N}$, it follows that $|[0, \nu]| \geq 2^{\aleph_0}$. Consequently, the cardinality of the ν distinct $f(\rho_k)$ is greater than or equal to $2^{\aleph_0}$. Since $i[\mathcal{W}]$ is denumerable, the cardinality of the set of purely subtle alphabet symbols contained in the set of ν distinct $f(\rho_k) \geq 2^{\aleph_0}$. This complete the proof. ■

With respect to the proof of Theorem 7.3.2, the function f determines the alphabet composition of the ultraword w . The word w is unreadable not only due to its infinite length but also due to the fact that it is composed of infinitely many purely subtle alphabet symbols.

The developmental paradigm d utilized for the two previous theorems is composed entirely of readable sentences. We now investigate what happens if a developmental paradigm contains countably many unreadable sentences. Let the nonempty developmental paradigm d' be composed of at most countably many members of ${}^*\mathcal{E} - \mathcal{E}$ and let $d' \subset {}^*\mathbf{B} \subset {}^*\mathbf{P}_0$. Construct, as previously, the set M_B from $\mathbf{B}$, rather than from $\mathbf{d}$ and suppose that $B \cap M_i = \emptyset$, $i \neq 0$. [This last requirement for B can be achieved as follows: construct a special symbol not originally in $\mathcal{A}$. Then this symbol along with $\mathcal{A}$ is considered the alphabet. Next only consider a B that does not contain this special symbol within any of its members. Now use this special symbol consistently with or without the spacing symbol to construct M_i $i \neq 0$. Of course $\wedge$ is interpreted as this special symbol with or without the spacing symbol in the axiom system S .]

The w that exists by the next theorem generates, upon applying the ultralogic ${}^*\mathbf{S}$, any selected event sequence that contains not only previously selected Natural events but also selected UN-events.

Theorem 7.3.3 *Let $d' = \{[g_i] \mid i \in \mathbb{N}\}$. Then there exists an ultraword $w \in {}^*\mathbf{M}_B - {}^*\mathbf{B}$ such that for each $i \in \mathbb{N}$, $[g_i] \in {}^*\mathbf{S}(\{w\})$.*

Proof. Consider the internal binary relation $G = \{(x, y) \mid (x \in {}^*\mathbf{B}) \wedge (y \in {}^*\mathbf{M}_B - {}^*\mathbf{B}) \wedge (x \in {}^*\mathbf{S}(\{y\}))\}$. Note that members of d' are members of ${}^\sigma\mathcal{E}$ or, at the most, denumerably many members of ${}^*\mathcal{E} - {}^\sigma\mathcal{E}$. From the analysis in the proof of Theorem 7.3.1, for a finite $F \subset \mathbf{B}$, there exists some $y \in \mathbf{M}_B - \mathbf{B}$ such that $F \subset S(\{y\})$. It follows by $*$ -transfer that if F is a finite or $*$ -finite subset of ${}^*\mathbf{B}$, then there exists some $y \in {}^*\mathbf{M}_B - {}^*\mathbf{B}$ such that $F \subset {}^*\mathbf{S}(\{y\})$. As in the proof of Theorem 7.3.1, this yields that G is at least concurrent on ${}^*\mathbf{B}$. However, $d' \subset {}^*\mathbf{B}$ and $|d'| \leq \aleph_0$. From $\aleph_1$ -saturation, there exists some $w \in {}^*\mathbf{M}_B - {}^*\mathbf{B}$ such that for each $[g_i] \in d'$, $[g_i] \in {}^*\mathbf{S}(\{w\})$. This completes the proof. ■

Let $\emptyset \neq \lambda \subset \mathbb{N}$ and $\mathcal{D}_j = \{d_{ij} \mid i \in \lambda\}$ and each $d_i \subset {}^*\mathbf{B}$ is considered to be a developmental paradigm either of type d or type d' and $B \cap M_i = \emptyset$, $i \neq 0$. Notice that $\mathcal{D}_j$ may be either a finite or denumerable set and Theorem 7.3.1 holds for the case that $\mathbf{d} \subset \mathbf{B}$, where $w \in {}^*\mathbf{M}_B - {}^*\mathbf{B}$. For each $d_{ij} \in \mathcal{D}_j$, use the Axiom of Choice to select an ultraword $w_{ij} \in {}^*\mathbf{M}_B - {}^*\mathbf{B}$ that exists by Theorems 7.3.1 (extended) and 7.3.3. Let $\{w_{ij} \mid i \in \lambda\}$ be such a set of ultrawords.

This next theorem shows the existence of an ultimate ultraword that by application of the ultralogic yields all of the other needed ultrawords. Thus w'_i generates, after application of the ultralogic $^*\mathbf{S}$, a Natural-system's event sequence, whether it contains only Natural events or selected UN-events. Then w' and $^*\mathbf{S}(\{w'\})$ solve Wheeler's "general grand unification problem."

Theorem 7.3.4 *There exists an ultimate ultraword $w'_j \in {}^*\mathbf{M}_B - {}^*\mathbf{B}$ such that for each $i \in \lambda$, $w_{ij} \in {}^*\mathbf{S}(\{w'_j\})$ and, hence, for each $d_{ij} \in \mathcal{D}_j$, $d_{ij} \subset {}^*\mathbf{S}(\{w_{ij}\}) \subset {}^*\mathbf{S}(\{w'_j\})$.*

Proof. For each finite $\{F_1, \dots, F_n\} \subset \mathbf{M}_B - \mathbf{B}$ there is a natural number, say m , such that for $i = 1, \dots, n$, $F_i \in M_j$ for some $j \leq m$. Hence, taking $b \in M_{m+1}$, we obtain that each $F_i \in S(\{b\})$. Observe that $b \notin \mathbf{B}$. By * -transfer, it follows that the internal relation $G = \{(x, y) \mid (x \in {}^*\mathbf{M}_B - {}^*\mathbf{B}) \wedge (y \in {}^*\mathbf{M}_B - {}^*\mathbf{B}) \wedge (x \in {}^*\mathbf{S}(\{y\}))\}$ is concurrent on internal ${}^*\mathbf{M}_B - {}^*\mathbf{B}$ and $\{w_{ij} \mid i \in \lambda\} \subset {}^*\mathbf{M}_B - {}^*\mathbf{B}$. Again $\aleph_1$ -saturation yields that there is some $w'_j \in {}^*\mathbf{M}_B - {}^*\mathbf{B}$ such that for each $i \in \lambda$, $w_{ij} \in {}^*\mathbf{S}(\{w'_j\})$. The last property is obtained from $d_{ij} \subset {}^*\mathbf{S}(\{w_{ij}\}) \subset {}^*\mathbf{S}({}^*\mathbf{S}(\{w'_j\})) = {}^*\mathbf{S}(\{w'_j\})$ since $\{w_i\}$ is an internal subset of ${}^*\mathbf{P}_0$. This completes the proof. ■

Corollary 7.3.4.1 *Let $\emptyset \neq \gamma \subset \mathbb{N}$. There exists an ultimate ultraword $w' \in {}^*\mathbf{M}_B - {}^*\mathbf{B}$ such that for each $j \in \gamma$, $w'_j \in {}^*\mathbf{S}(\{w'\})$ and, hence, for each $d_{ij} \in \bigcup \mathcal{D}_j$, $d_{ij} \subset {}^*\mathbf{S}(\{w'_j\}) \subset {}^*\mathbf{S}(\{w'\})$.*

The same analysis used to obtain Theorem 7.3.2 can be applied to the ultrawords of Theorems 7.3.3 and 7.3.4.

7.4 Ultracontinuous Deduction

In 1968, a special topology on the set of all nonempty subsets of a given set X was constructed and investigated by your author. We apply a similar topology to subsets of $\mathcal{E}$.

Suppose that nonempty $X \subset \mathcal{E}$. Let τ be the discrete topology on X . In order to topologize $\mathcal{P}(X)$, proceed as follows: for each $G \in \tau$, let $N(G) = \{A \mid (A \subset X) \wedge (A \subset G)\} = \mathcal{P}(G)$. Consider $\mathcal{B} = \{N(G) \mid G \in \tau\}$ to be a base for a topology τ_1 on $\mathcal{P}(X)$. Let $A \in N(G_1) \cap N(G_2)$. The discrete topology implies that $N(A)$ is a base element and that $N(A) \subset N(G_1) \cap N(G_2)$. There is only one member of $\mathcal{B}$ that contains X and this is $\mathcal{P}(X)$. Thus if $\mathcal{P}(X)$ is covered by members of $\mathcal{B}$, then $N(X) = \mathcal{P}(X)$ is one of these covering objects. Thus $(\mathcal{P}(X), \tau_1)$ is a compact space. Further, since $N(\emptyset) \subset N(G)$ for each $G \in \tau$, the space $(\mathcal{P}(X), \tau_1)$ is connected. The topology τ_1 is a special case of a more general topology with the same properties. [2] Suppose that $D \subset X$. Let $D \in N(G) = \mathcal{P}(G)$, $G \in \tau$. Then $D \in N(D) \subset N(G)$. This yields that the nonstandard monad is $\mu(D) = \bigcap \{^*N(G) \mid N(G) \in \mathcal{B}\} = {}^*(\mathcal{P}(D)) = {}^*\mathcal{P}({}^*D)$.

This next theorem shows that the ultralogic $^*\mathbf{S}$ produces Natural and ultranatural events in an ultracontinuous manner.

Theorem 7.4.1 *Any consequence operator $C: (\mathcal{P}(X), \tau_1) \rightarrow (\mathcal{P}(X), \tau_1)$ is continuous.*

Proof. Let $A \in \mathcal{P}(X)$ and $H \in {}^*\mathbf{C}[\mu(A)]$. Then there exists some $B \in \mu(A)$ such that ${}^*\mathbf{C}(B) = H$. Hence, $B \in {}^*\mathcal{P}({}^*A)$. By * -transfer of a basic property of our consequence operators, ${}^*\mathbf{C}(B) \subset {}^*\mathbf{C}({}^*A) = {}^*(\mathbf{C}(A))$. Thus ${}^*(\mathbf{C}(B)) \in {}^*(\mathcal{P}(\mathbf{C}(A)))$ implies that ${}^*\mathbf{C}(B) \in \mu(A)$. Therefore, ${}^*\mathbf{C}[\mu(A)] \subset \mu[\mathbf{C}(A)]$. Consequently, $\mathbf{C}$ is continuous. ■

Corollary 7.4.1.1 *For any $X \subset \mathcal{E}$, and any consequence operator $\mathbf{C}: \mathcal{P}(X) \rightarrow \mathcal{P}(X)$, the map ${}^*\mathbf{C}: {}^*(\mathcal{P}(X)) \rightarrow {}^*(\mathcal{P}(X))$ is ultracontinuous.*

Corollary 7.4.1.2 *Let d [resp. d' , d or d'] be a developmental paradigm as defined for Theorem 7.3.1 [resp. Theorem 7.3.3, 7.3.4]. Let w be a ultraword that exists by Theorem 7.3.1 [resp Theorem 7.3.3, 7.3.4]. Then d [resp. d' , d or d'] is obtained by means of a ultracontinuous subtle deductive process applied to $\{w\}$.*

Recall that in the real valued case, a function $f: [a, b] \rightarrow \mathbb{R}$ is uniformly continuous on $[a, b]$ iff for each $p, q \in {}^*[a, b]$ such that $p - q \in \mu(0)$, then $f(p) - f(q) \in \mu(0)$. If $D \subset [a, b]$ is compact, then $p, q \in {}^*D$ and $p - q \in \mu(0)$ imply that there is a standard $r \in D$ such that $p, q \in \mu(r)$. Also, for each $r \in D$ and any $p, q \in \mu(r)$, it follows that $p - q \in \mu(r)$. Thus, if compact $D \subset [a, b]$, then $f: D \rightarrow \mathbb{R}$ is uniformly continuous iff for every $r \in D$ and each $p, q \in \mu(r)$, ${}^*f(p), {}^*f(q) \in \mu(f(r))$. With this characterization in mind, it is clear that any consequence operator $\mathbf{C}: \mathcal{P}(X) \rightarrow \mathcal{P}(X)$ satisfies the following statement. For each $A \in \mathcal{P}(X)$ and each $p, q \in \mu(A)$, ${}^*\mathbf{C}(p), {}^*\mathbf{C}(q) \in \mu(\mathbf{C}(A))$.

From the above discussion, one can think of ultracontinuity as being a type of ultrauniform continuity.

7.5 Hypercontinuous Gluing

There are various methods that can be used to investigate the behavior of adjacent frozen segments. All of these methods depend upon a significant result relative to discrete real or vector valued functions. The major goal in this section is to present a complete proof of this major result and to indicate how it is applied.

First, as our standard structure, consider either the intuitive real numbers as atoms or axiomatically a standard structure with atoms $\mathbf{ZFR} = \mathbf{ZF} + \mathbf{AC} + A_1(\text{atoms}) + A(\text{atoms}) + |A| = c$, where A is isomorphic to the real numbers and $A_1 \cap A = \emptyset$. Then, as done previously, there is a model $\langle C, \in, = \rangle$ within our $\mathbf{ZF} + \mathbf{AC}$ model for $\mathbf{ZFR}$, where A has all of the ordered field properties as the real numbers. A superstructure $\langle \mathcal{R}, \in, = \rangle$ is constructed in the usual manner, where the superstructure $\langle \mathcal{N}, \in, = \rangle$ is a substructure. Proceeding as in Chapter 2, construct ${}^*\mathcal{M}_1 = \langle {}^*\mathcal{R}, \in, = \rangle$ and $\mathcal{Y}_1$. The structure $\mathcal{Y}_1$ is called the *Extended Grundlegend Structure* — the EGS. The Grundlegend Structure is a substructure of $\mathcal{Y}_1$.

It is important to realized in what follows that the objects utilized for the G-structure *interpretations* are nonempty finite equivalence classes of partial sequences. Due to this fact, the following results should not lead to ambiguous interpretations.

As a preliminary to the technical aspects of this final section, we introduce the following definition. A function $f: [a, b] \rightarrow \mathbb{R}^m$ is *differentiable-C* on $[a, b]$ if it is continuously differentiable on (a, b) except at finitely many removable discontinuities. This definition is extended to the end points $\{a, b\}$ by application of one-sided derivatives. For any $[a, b]$, consider a partition $P = \{a_0, a_1, \dots, a_n, a_{n+1}\}$, $n \geq 1$, $a = a_0$, $b = a_{n+1}$ and $a_{j-1} < a_j$, $1 \leq j \leq n+1$. For any such partition P , let the real valued function g be defined on the set $D = [a_0, a_1] \cup (a_1, a_2) \cup \dots \cup (a_n, a_{n+1}]$ as follows: for each $x \in [a_0, a_1]$, let $g(x) = r_1 \in \mathbb{R}$; for each $x \in (a_{j-1}, a_j]$, let $g(x) = r_j \in \mathbb{R}$, $1 < j \leq n$; for each $x \in (a_n, b]$, let $g(x) = r_{n+1} \in \mathbb{R}$. It is obvious that g is a type of simple step function. Notationally, let $\mathcal{F}(A, B)$ denote the set of all functions with domain A and codomain B .

Theorem 7.5.1. shows that if a time fracture occurs of the minimum or intermediate type, then there exists an ultracontinuous, ultrauniform and ultrasmooth alteration process within the NSP-world that yields all of the alterations.

Theorem 7.5.1 *There exists a function $G \in {}^*(\mathcal{F}([a, b], \mathbb{R}))$ with the following properties.*

- (i) *The function G is * -continuously * -differentiable and * -uniformly * -continuous on ${}^*[a, b]$,*
- (ii) *for each odd $n \in {}^*\mathbb{N}$, ($n \geq 3$), G is * -differentiable- C of order n on ${}^*[a, b]$,*
- (iii) *for each even $n \in {}^*\mathbb{N}$, G is * -continuously * -differentiable in ${}^*[a, b]$ except at finitely many points,*
- (iv) *if $c = \min\{r_1, \dots, r_{n+1}\}$, $d = \max\{r_1, \dots, r_{n+1}\}$, then the range of $G = {}^*[c, d]$, $\mathbf{st}(G)$ at least maps D into $[c, d]$ and $(\mathbf{st}(G))|D = g$.*

Proof. First, for any real c, d , where $d \neq 0$, consider the finite set of functions

$$(7.5.1) \quad h_j(x, c, d) = (1/2)(r_{j+1} - r_j) \left(\sin((x - c)\pi/(2d)) + 1 \right) + r_j,$$

$1 \leq j \leq n$. Each h_j is continuously differentiable for any order at each $x \in \mathbb{R}$. Observe that for each odd $m \in \mathbb{N}$, each m 'th derivative $h_j^{(m)}$ is continuous at $(c + d)$ and $(c - d)$ and $h_j^{(m)}(c + d) = h_j^{(m)}(c - d) = 0$ for each j .

Let positive $\delta \in \mu(0)$. Consider the finite set of internal intervals $\{[a_0, a_1 - \delta), (a_1 + \delta, a_2 - \delta), \dots, (a_n + \delta, b]\}$ obtained from the partition P . Denote these intervals in the expressed order by I_j , $1 \leq j \leq n + 1$. Define the internal function

$$(7.5.2) \quad G_1 = \{(x, r_1) | x \in I_1\} \cup \dots \cup \{(x, r_{n+1}) | x \in I_{n+1}\}.$$

Let internal $I_j^\dagger = [a_j - \delta, a_j + \delta]$, $1 \leq j \leq n$, and for each $x \in I_j^\dagger$, let internal

$$(7.5.3) \quad G_j(x) = (1/2)(r_{j+1} - r_j) \left({}^*\sin((x - a_j)\pi/(2\delta)) + 1 \right) + r_j.$$

Define the internal function

$$(7.5.4) \quad G_2 = \{(x, G_1(x)) | x \in I_1^\dagger\} \cup \dots \cup \{(x, G_n(x)) | x \in I_n^\dagger\}.$$

The final step is to define $G = G_1 \cup G_2$. Then $G \in {}^*(\mathcal{F}([a, b], \mathbb{R}))$.

By * -transfer, the function G_1 has an internal * -continuous * -derivative $G_1^{(1)}$ such that $G_1^{(1)}(x) = 0$ for each $x \in I_1 \cup \dots \cup I_{n+1}$. Applying * -transfer to the properties of the functions $h_j(x, c, d)$, it follows that G_2 has a unique internal * -continuous * -derivative

$$(7.5.5) \quad G_2^{(1)} = (1/(4\delta))(r_{j+1} - r_j)\pi \left({}^*\cos((x - a_j)\pi/(2\delta)) \right)$$

for each $x \in I_1^\dagger \cup \dots \cup I_n^\dagger$. The results that the * -left limit for the internal $G_1^{(1)}$ and the * -right limit for internal $G_2^{(1)}$ at each $a_j - \delta$ as well as the * -left limit of $G_2^{(1)}$ and * -right limit of $G_1^{(1)}$ at each $a_j + \delta$ are equal to 0 and $0 = G_2^{(1)}(a_j - \delta) = G_2^{(1)}(a_j + \delta)$ imply that internal G has a * -continuous * -derivative $G^{(1)} = G_1^{(1)} \cup G_2^{(1)}$ defined on ${}^*[a, b]$.

A similar analysis and * -transfer yield that for each $m \in {}^*\mathbb{N}$, $m \geq 2$, G has an internal * -continuous * -derivative $G^{(m)}$ defined at each $x \in {}^*[a, b]$ except at the points $a_j \pm \delta$ whenever $r_{j+1} \neq r_j$. However, it is obvious from the definition of the functions h_j that for each odd $m \in {}^*\mathbb{N}$, $m \geq 3$, each internal $G^{(m)}$ can be made * -continuous at each $a_j \pm \delta$ by simply defining $G^{(m)}(a_j \pm \delta) = 0$ and with this parts (i), (ii), and (iii) are established.

For part (iv), assume that $r_j \leq r_{j+1}$. From the definition of the functions h_j , it follows that for each $x \in I_j \cup I_j^\dagger \cup I_{j+1}$, $r_j \leq G(x) \leq r_{j+1}$. The nonstandard intermediate value theorem

implies that $G[*[a_j, a_{j+1}]] = [*[r_j, r_{j+1}]]$ and in like manner for the case that $r_{j+1} < r_j$. Hence, $G[*[a, b]] = [*[c, d]]$. Clearly, $\text{st}(*D) = [a, b]$. If $p \in D$ and $x \in \mu(p) \cap *D$, then $G(x) = r_j = g(p)$ for some j such that $1 \leq j \leq n+1$. This completes the proof. ■

The nonstandard approximation theorem 7.5.1 can be extended easily to functions that map D into $\mathbb{R}^m$. For example, assume that $F: D \rightarrow \mathbb{R}^3$, the component functions F_1, F_2 are continuously differentiable on $[a, b]$; but that F_3 is a g type step function on D . Then letting $H = (*F_1, *F_2, G)$, on $*[a, b]$, where G is defined in Theorem 4.1, we have an internal $*$ -continuously $*$ -differentiable function $H: *[a, b] \rightarrow *\mathbb{R}^3$, with the property that $\text{st}(H)|D = F$.

With respect to Theorem 7.5.1, it is interesting to note that if h_j is defined on $\mathbb{R}$, then for even orders $n \in \mathbb{N}$,

$$|h_j^{(n)}(c \pm d)| = \left| \frac{(r_{j+1} - r_j)\pi^n}{2^{n+1}d^n} \right| = 0 \quad (7.5.6)$$

for $r_{j+1} = r_j$ but not 0 otherwise. If $r_{j+1} - r_j \neq 0$, then $G_2^{(n)}(a_j \pm \delta)$ is an infinite nonstandard real number. Indeed, if m_i is an increasing sequence of even numbers in $*\mathbb{N}$ and $r_{j+1} \neq r_j$, then $|G_2^{(m_i)}(a_j \pm \delta)|$ forms a decreasing sequence of nonstandard infinite numbers. The next result is obvious from the previous result.

Corollary 7.5.1.1 *For each $n \in *\mathbb{N}$, then internal $G^{(n)} = G_1^{(n)} \cup G_2^{(n)}$ is $*$ -bounded on $*[a, b]$.*

Let $D(a, b)$ be the set of all bounded and piecewise continuously differentiable functions defined on $[a, b]$. By considering all of the possible (finitely many) subintervals, where $f \in D(a, b)$, it follows from the Riemann sum approach that for each real $\nu > 0$, there exists a real $\nu_1 > 0$ such that for each real ν_i , $0 < \nu_i < \nu_1$, a sequence of partitions $P_i = \{a = b_0^i < \dots < b_{k_i}^i = b\}$ can be selected such that the $\text{mesh}(P_i) \leq \nu_i$ and

$$|(f(b) - f(a)) - \sum_{n=1}^{k_i} f'(t_n)(b_n^i - b_{n-1}^i)| < \nu \quad (7.5.7)$$

for any $t_n \in (b_{n-1}^i, b_n^i)$.

Moreover, for any given number M , the sequence of partitions can be so constructed such that there exists a j such that for each $i > j$, $k_i > M$, where P_i and P_j are partitions within the sequence of partitions. By $*$ -transfer of these facts and by application of Theorem 7.5.1 and its corollary we have the next result.

Corollary 7.5.1.2 *For each $n \in \mathbb{N}$ and each internal $G^{(n)}$, the difference $G^{(n)}(b) - G^{(n)}(a)$ is infinitesimally close to an (externally) infinity $*$ -finite sum of infinitesimals.*

A developmental paradigm is a very general object and, therefore, can be used for numerous applications. At present, developmental paradigms are still being viewed from the *substratum* or external world. For what follows, it is assumed that a developmental paradigm d traces the evolutionary history of a specifically named natural system or systems. In this first application, let each $F_i \in d$ have the following property (**P**).

F_i describes “the general behavior and characteristics of the named natural system S_1 as well as the behavior and characteristics of named constituents contained within S_1 at time t_i .”

Recall that for $F_i, F_{i+1} \in d$, there exist unique functions $f_0 \in \mathbf{F}_i = [f]$, $g_0 \in \mathbf{F}_{i+1} = [g]$ such that $f_0, g_0 \in T^0$ and $\{(0, f_0(0))\} \in [f]$, $\{(0, g_0(0))\} \in [g]$. Thus, to each $F_j \in d$, correspond the

unique natural number $f_0(0)$. Let $D = [t_{i-1}, t_i) \cup (t_i, t_{i+1}]$ and define $f_1: D \rightarrow \mathbb{N}$ as follows: for each $x \in [t_{i-1}, t_i)$, let $f_1(x) = f_0(0)$; for each $x \in (t_i, t_{i+1}]$, let $f_1(x) = g_0(0)$. Application of theorem 7.5.1 yields the internal function G such that $G|D = f_1$. For these physical applications, utilize the term “substratum” in the place of the technical terms “pure nonstandard.” [Note: Of course, elsewhere, the term “pure NSP-world” or simply the “NSP-world” is used as a specific name for what has here been declared as the substratum.] This yields the following statements, where the symbols F_i and F_{i+1} are defined and characterized by the expression inside the quotation marks in property **(P)**.

(A): There exists a substratum hypercontinuous, hypersmooth, hyperuniform process G that binds together F_i and F_{i+1} .

(B): There exists a substratum hypercontinuous, hypersmooth, hyperuniform alteration process G that transforms F_i into F_{i+1} .

(C): There exists an ultracontinuous subtle force-like (i.e. deductive) process that yields F_i for each time t_i within the development of the natural system.

In order to justify **(A)** and **(B)**, specific measures of physical properties associated with constituents may be coupled together. Assume that for a subword $r_i \in F_i \in \mathbf{d}$, the symbols r_i denote a numerical quantity that aids in characterizing the behavior of an object in a system S_1 or the system itself. Let **(M₁)** be the statement:

“There exists a substratum hypercontinuous, hypersmooth, hyperuniform functional process G_i such that G_i when restricted to the standard mathematical domain it is f_i and such that G_i hypercontinuously changes r_i for system S_1 at time t_i into r_{i+1} for system S_1 at time t_{i+1} .”

This modeling procedure yields the following interpretation:

(D) If there exists a continuous or uniform [resp. discrete] functional process f_i that changes r_i for S_1 at time t_i into r_{i+1} for S_1 at time t_{i+1} , then **(M₁)**.

At a particular moment t_i , two natural systems S_1 and S_2 may interface. More generally, two very distinct developmental paradigms may exist one $\mathbf{d}_1$ at times prior to t_i (in the t_i past) and one $\mathbf{d}_2$ at time after t_i (in the t_i future). We might refer to the time t_i as a *standard time fracture*. Consider the developmental paradigm $\mathbf{d}_3 = \mathbf{d}_1 \cup \mathbf{d}_2$. In this case, the paradigms may be either of type $\mathbf{d}$ or $\mathbf{d}'$. For the type $\mathbf{d}'$, the corresponding system need not be considered a natural system but could be a pure substratum system.

At t_i an $F_i \in \mathbf{d}_3$ can be characterized by statement **(P)** (with the term natural removed if F_i is a member of a $\mathbf{d}'$). In like manner, F_{i+1} at time t_{i+1} can be characterized by **(P)**. Statements **(A)**, **(B)**, **(C)** can now be applied to $\mathbf{d}_3$ and a modified statement **(D)**, where the second symbol string S_1 is changed to S_2 . Notice that this modeling applies to the actual human ability that only allows for two discrete descriptions to be given, one for the interval $[t_{i-1}, t_i)$ and one for the interval $(t_i, t_{i+1}]$. From the modeling viewpoint, this is often sufficient since the length of the time intervals can be made smaller than Planck time.

Recall that an analysis of the scientific method used in the investigation of natural system should take place exterior to the language used to describe the specific system development. Suppose that $\mathcal{D}$ is the language accepted for a scientific discipline and that within $\mathcal{D}$ various expressions from mathematical theories are used. Further, suppose that enough of the modern theory of sets is employed so that the EGS can be constructed. The following statement would hold true for $\mathcal{D}$.

If by application of first-order logic to a set of non-mathematical premises taken from $\mathcal{D}$ it is claimed that it is not logically possible for statements such as (A), (B), (C) and (D) to hold, then the set of premises is inconsistent.

CHAPTER 7 REFERENCES

- 1 Beltrametti, E. G. and G. Cassinelli, The logic of quantum mechanics, in *Encyclopedia of Mathematics and its Application*, Vol. 15, Addison-Wesley, Reading, 1981.
- 2 Herrmann, R. A., *Some Characteristics of Topologies on Subsets of a Power Set*, University Microfilm, M-1469, 1968.
- 3 Herrmann, R. A., *Einstein Corrected*, (1993) I. M. P. <http://www.arXiv.org/abs/math/0312189>

NOTES

- [1] The actual members, F_i , of a developmental paradigm d need not be unique. However, the specific information contained in each readable word used for a specific $i \in \mathbb{N}$ is unique.
- [2] Depending upon the application, a single standard word may also be termed as an ultraword.
- [3] For a new more detailed method to obtain ultrawords for a refined developmental paradigm, see pages 4 – 7 of <http://arxiv.org/abs/math/0605120>

.....

9.1 An Extension.

Although it is often not necessary, we assume when its useful that we are working within the EGS. Further, this structure is assumed to be $2^{|\mathcal{M}_1|}$ -saturated, where $\mathcal{M}_1 = \langle \mathcal{R}, \in, = \rangle$, even though such a degree of saturation can usually be reduced. Referring to the paragraph prior to Theorem 7.3.3, it can be assumed that the developmental paradigm $d' \subset {}^*\mathbf{B} \subset {}^*\mathbf{P}_0$. It is not assumed that such a developmental paradigm is obtained from the process discussed in Theorem 7.2.1, although a modification of the proof of Theorem 7.2.1 appears possible in order to allow this method of selection.

Theorem 9.1.1 *Let $d' = \{[g_i] \mid i \in \lambda\}$, $|\lambda| < 2^{|\mathcal{M}_1|}$. There exists an ultraword $w \in {}^*\mathbf{M}_B - {}^*\mathbf{B}$ such that for each $i \in \lambda$, $[g_i] \in {}^*\mathbf{S}(\{w\})$.*

Proof. The same as Theorem 7.3.3 with the change in saturation. ■

Let $\mathcal{D} = \{d_i \mid i \in \lambda\}$, $|\lambda| < 2^{|\mathcal{M}_1|}$, $|d_i| < 2^{|\mathcal{M}_1|}$ and each $d_i \subset {}^*\mathbf{B}$ is considered to be a developmental paradigm either of type d or type d' . For each $d_i \in \mathcal{D}$, use the Axiom of Choice to select an ultraword $w_i \in {}^*\mathbf{M}_B - {}^*\mathbf{B}$ that exists by Theorems 9.1.1. Let $\{w_i \mid i \in \lambda\}$ be such a set of ultrawords.

Theorem 9.1.2 *There exists an ultraword $w' \in {}^*\mathbf{M}_B - {}^*\mathbf{B}$ such that for each $i \in \lambda$, $w_i \in {}^*\mathbf{S}(\{w'\})$ and, hence, for each $d_i \in \mathcal{D}$, $d_i \subset {}^*\mathbf{S}(\{w'\})$.*

Proof. The same as Theorem 7.3.4 with the change in saturation. ■

9.2 NSP-World Alphabets.

First, recall the following definition. $P_m = \{f \mid (f \in T^m) \wedge (\exists z((z \in \mathcal{E}) \wedge (f \in z) \wedge \forall x((x \in \mathbb{N}) \wedge (x > m) \rightarrow \neg \exists y((y \in T^x) \wedge (y \in z))))))\}$. The set $T = i[\mathcal{W}]$. The set P_m determines the unique

partial sequence $f \in [g] \in \mathcal{E}$ that yields, for each $j \in \mathbb{N}$ such that $0 \leq j \leq m$, $f(j) = i(a)$, where $i(a)$ is an “encoding” in A_1 of the alphabet symbol “a” used to construct our intuitive language $\mathcal{W}$. The set $[g]$ represents an intuitive word constructed from such an alphabet of symbols.

Within the discipline of Mathematical Logic, it is assumed that there exists symbols — a sequence of variables — each one of which corresponds, in a one-to-one manner, to a natural number. Further, under the subject matter of generalized first-order theories [2], it is assumed that the cardinality of the set of constants is greater than $\aleph_0$. In the forthcoming investigation, it may be useful to consider an alphabet that injectively corresponds to the real numbers $\mathbb{R}$. This yields a new alphabet $\mathcal{A}'$ containing our original alphabet. A new collection of words $\mathcal{W}'$ composed of nonempty finite strings of such alphabet symbols may be constructed. It may also be useful to well-order $\mathbb{R}$. The set $\mathcal{E}$ also exists with respect to the set of words $\mathcal{W}'$. Using the ESG, many previous results in this book now hold with respect to $\mathcal{W}'$ and for the case that we are working in a $2^{|\mathcal{M}_1|}$ -saturated enlargement.

With respect to this extended language, if you wish to except the possibility, a definition as to what constitutes a purely subtle alphabet symbol would need to be altered in the obvious fashion. Indeed, for T in the definition of P_m , we need to substitute $T' = i[\mathcal{W}']$. Then the altered definition would read that $r \in {}^*A_1 \simeq {}^*\mathbb{R}$ is a *pure subtle alphabet symbol* if there exists an $m \in \mathbb{N}$ and $f \in {}^*(P_m)$, or if $m \in {}^*\mathbb{N} - \mathbb{N}$ and $f \in P_m$, and some $j \in {}^*\mathbb{R}$ such that $f(j) = r \notin i[\mathcal{W}']$. Further, some of the previous theorems also hold when the proofs are modified.

Although these extended languages are of interest to the mathematician, most of science is content with approximating a real number by means of a rational number. In all that follows, the cardinality of our language, if not denumerable, will be specified. All theorems from this book that are used to establish a result relative to a denumerable language will be stated without qualification. If a theorem has not been reestablished for a higher language but can be so reestablished, then the theorem will be termed an *extended* theorem.

9.3 General Paradigms.

There is the developmental paradigm, and for nondetailed descriptions the *general developmental paradigm*. But now we have something totally new — the *general paradigm*. It is important to note that the general paradigm is considered to be distinct from developmental paradigms, although certain results that hold for general paradigms will hold for developmental paradigms and conversely. For example, associated with each general paradigm G_A is an ultraword w_g such that the set $G_A \subset {}^*\mathbf{S}(\{w_g\})$ and all other theorems relative to such ultrawords hold for general paradigms. The general paradigm is a collection of words that discuss, in general, the behavior of entities and other constituents of a natural system. They, usually, do not contain a time statement W_i as it appears in section 7.1 for developmental paradigm descriptions. Our interest in this section is relative to only two such general paradigms. The reader can easily generate many other general paradigms.

Let c' be a symbol that denotes some fixed real number and n' a symbol that denotes a natural number. [Note: what follows is easily extended to an extended language.] Suppose that you have a theory which includes each member of the following set (i suppressed).

$$(9.3.1) \quad G_A = \{An||\text{elementary}||\text{particle}||\alpha(n')||\text{with}||\text{kinetic}||\text{energy}||c'+1/(n'). \mid n \in G \wedge n \neq 0\},$$

where G is a denumerable subset of the real numbers. The set G_A is of particular interest when $G = \mathbb{N}$. Theories that include such sentences consider such particles as *free in space*.

Theorem 9.3.1 shows that it is necessary for subparticles to exist.

Of particular interest is the composition of members of ${}^*\mathbf{G}_A - \mathbf{G}_A$. Notice that $|\mathbf{G}_A| = |G|$ since $z_1, z_2 \in \mathbf{G}_A$ and $z_1 \neq z_2$ iff $[x_1] = z_1, [x_2] = z_2, x_1(30) = x_1(2) \neq x_2(30) = x_2(2), x_1(2), x_2(2) \in G$. Now consider the bijection $K: \mathbf{G}_A \rightarrow G$.

Theorem 9.3.1 *The set $[g] \in {}^*\mathbf{G}_A - \mathbf{G}_A$ iff there exists a unique $f \in {}^*(P_{55})$ and $\nu \in {}^*G - G$ such that $[g] = [f]$, and $f(55) = i(A), f(54) = i(n), f(53) = i(|||), \dots, f(30) = f(2), \dots, f(3) = i(), f(2) = \nu \in {}^*G - G \subset {}^*\mathbb{R} - \mathbb{R}, f(1) = i(), f(0) = ()$.*

Proof. From the definition of $\mathbf{G}_A$ the sentence

$$\begin{aligned} \forall z(z \in \mathcal{E} \rightarrow ((z \in \mathbf{G}_A) \leftrightarrow \exists!x\exists!w((w \in G) \wedge (x \in P_{55}) \wedge (x \in z) \wedge \\ ((55, i(A)) \in x) \wedge ((54, i(n)) \in x) \wedge \dots \wedge (x(30) = x(2)) \wedge \dots \wedge \\ ((3, i()) \in x) \wedge (x(2) = w) \wedge (K(z) = w) \wedge \\ ((1, i()) \in x) \wedge ((0, i()) \in x))))). \end{aligned} \quad (9.3.2)$$

holds in $\mathcal{M}$, hence in ${}^*\mathcal{M}$. From the fact that K is a bijection, it follows that ${}^*K[{}^*\mathbf{G}_A - \mathbf{G}_A] = {}^*G - G \subset {}^*\mathbb{R} - \mathbb{R}$. The result now follows from * -transfer. ■

Using Theorem 9.3.1, each member of ${}^*\mathbf{G}_A - \mathbf{G}_A$, when interpreted by considering i^{-1} , has only two positions with a single missing object since positions 30 and 2 do not correspond to any symbol string in our language $\mathcal{W}$. This interpretation still retains a vast amount of content, however. For a specific member, you could substitute a new constructed symbol, not in $\mathcal{W}$, into these two missing positions. Depending upon what type of pure nonstandard number this inserted symbol represents, the content of such a sentence could be startling. Let Γ' be a nonempty set of new symbols disjoint from $\mathcal{W}$ and assume that Γ' is injectively mapped by H into ${}^*G - G$.

Although human ability may preclude the actual construction of more than denumerably many new symbols, you might consider this mapping to be onto if you accept the ideas of extended languages with a greater cardinality. As previously, denote these new symbols by γ' . Now let

$$(9.3.3) \quad \begin{aligned} G'_A = \{ \text{An} || \text{elementary} || \text{particle} || \alpha(\gamma') || \text{with} || \\ \text{kinetic} || \text{energy} || c' + 1/(\gamma') \cdot | H(\gamma') \in {}^*G - G \}, \end{aligned}$$

This leads to the following interpretation stated in terms of describing sets.

- (1) *The describing set $\mathbf{G}_A$ (mathematically) exists iff the describing set G'_A (mathematically) exists.*

9.4 Interpretations

Recall that the Natural world portion of the NSP-world model may contain *undetectable* objects, where “undetectable” means that there does not appear to exist human, or humanly constructible machine sensors that directly detect the objects or directly measure any of the objects physical properties. The rules of the scientific method utilized within the micro-world of subatomic physics allow all such undetectable Natural objects to be accepted as existing reality.[1] The properties of such objects are indirectly deduced from the observed properties of gross matter. In order to have indirect evidence of the objectively real existence of such objects, such indirectly obtained behavior will usually satisfy a specifically accepted model.

Although the numerical quantities associated with these undetectable Natural (i.e. standard) world objects, if they really do exist, cannot be directly measured, these quantities are still represented by standard mathematical entities. By the rules of correspondence for interpreting pure NSP-world entities, the members of G'_A must be considered as undetectable pure NSP-world objects, assuming any of them exist in this background world. On the other hand, when viewed within the EGS, any finite as well as many infinite subsets of G'_A are internal sets. Consequently, some finite collects of such objects *may be* assumed to indirectly effect behavior in the Natural world.

The concept of *realism* often dictates that all interpreted members of a mathematical model be considered as existing in reality. The philosophy of science that accepts only *partial realism* allows for the following technique. One can stop at any point within a mathematically generated physical interpretation. Then proceed from that point to deduce an intuitive physical theory, but only using other not interpreted mathematical formalism as auxiliary constructs or as catalysts. With respect to the NSP-world, another aspect of interpretation enters the picture. Assuming realism, then the question remains which, if any, of these NSP-entities actually indirectly influence Natural world processes? This interpretation process allows for the possibility that none of these pure NSP-world entities has any effect upon the standard world. These ideas should always be kept in mind.

If you accept that such particles as described by G_A can exist in reality, then the philosophy of realism leads to the next interpretation.

(2) *If there exist elementary particles with Natural-system behavior described by G_A , then there exist pure NSP-world objects that display within the NSP-world behavior described by members of G'_A .*

The concept of absolute realism would require that the acceptance of the elementary particles described by G_A is indirect evidence for the existence of the G'_A described objects. I caution the reader that the interpretation we apply to such sets of sentences as G_A are only to be applied to such sets of sentences.

The EGS may, of course, be interpreted in infinitely many different ways. Indeed, the NSP-world model with its physical-type language can also be applied in infinitely many ways to infinitely many scenarios. I have applied it to such models as the GGU-model, among others. In this section, I consider another possible interpretation relative to those Big Bang cosmologies that postulate real objects at or near infinite temperature, energy or pressure. These theories incorporate the concept of the *initial singularity(ies)*.

One of the great difficulties with many Big Bang cosmologies is that no meaningful physical interpretation for formation of the initial singularity is forthcoming from the theory itself. The fact that a proper and acceptable theory for creation of the universe requires that consideration not only be given to the moment of zero cosmic time but to what might have occurred “prior” to that moment in the nontime period is what partially influenced Wheeler to consider the concept of a *pregeometry*. [3], [4] It is totally unsatisfactory to dismiss such questions as “unmeaningful” simply because they cannot be discussed in your favorite theory. Scientists must search for a broader theory to include not only the question but a possible answer.

Although the initial singularity for a Big Bang type of state of affairs apparently cannot be discussed in a meaningful manner by many standard physical theories, it can be discussed by application of our NSP-world language. Let c' be a symbol that represents any fixed real number. Define

$$G_B = \{ \text{An} || \text{elementary} || \text{particle} || \alpha(n') || \text{with} || \text{total} || \text{energy} || c' + n'. \mid n \in \mathbb{N} \}, \quad (9.4.1)$$

Application of Theorem 9.3.1 to G_B yields the set

$$G'_B = \{ \text{An} || \text{elementary} || \text{particle} || \alpha(\gamma') || \text{with} || \text{total} || \text{energy} || c' + \gamma'. \mid \gamma \in {}^*\mathbb{N} - \mathbb{N} \}, \quad (9.4.2)$$

(3) *If there exist elementary particles with Natural-system behavior described by G_B , then there exist pure NSP-world objects that display within the NSP-world behavior described by members of G'_B .*

The particles being described by G'_B have various infinite energies. These infinite energies **do not** behave in the same manner as would the real number energy measures discussed in G_B . As is usual when a metalanguage physical theory is generated from a formalism, we can further extend and investigate the properties of the G'_B objects by imposing upon them the corresponding behavior of the positive infinite hyperreal numbers. This produces some interesting propositions. Hence, we are able to use a nonstandard physical world language in order to give further insight into the state of affairs at or near a cosmic initial singularity. This gives *one* solution to a portion of the pregeometry problem. I point out that there are other NSP-world models for the beginnings of our universe, if there was such a beginning. Of course, the statements in G'_B need not be related at all to any Natural world physical scenario, but could refer only to the behavior of pure NSP-world objects.

Notice that Theorems such as 7.3.1 and 7.3.4 relative to the generation of developmental paradigms by ultrawords, also apply to general paradigms, where M , M_B , P_0 are defined appropriately. The following is a slight extension of Theorem 7.3.2 for general paradigms. Theorem 9.4.1 will also hold for developmental paradigms.

Theorem 9.4.1 shows that there exists an ultraword w such that the basic ultralogic ${}^*\mathbf{S}$ when applied to w yields not only a specific collection of elementary particles but also a collection of subparticles. Further, it follows that there exists an ultraword w'' such that when ${}^*\mathbf{S}$ is applied to w'' the result is the collection all of the elementary particles that are claimed to exist and comprise the material portion of our universe.

Theorem 9.4.1 *Let G_C be any denumerable general paradigm. Then there exists an ultraword $w \in {}^*\mathbf{P}_0$ such that for each $\mathbf{F} \in G_C$, $\mathbf{F} \in {}^*\mathbf{S}(\{w\})$ and there exist infinitely many $[g] \in {}^*G_C - G_C$ such that $[g] \in {}^*\mathbf{S}(\{w\})$.*

Proof. In the proof of Theorem 7.3.2, it is shown that there exists some $\nu \in {}^*\mathbb{N} - \mathbb{N}$ such that ${}^*h[[0, \nu]] \subset {}^*\mathbf{S}(\{w\})$ and ${}^*h[[0, \nu]] \subset {}^*G_C$. Since $|{}^*h[[0, \nu]]| \geq 2^{\aleph_0}$, then $|{}^*h[[0, \nu]] - h[\mathbb{N}]| \geq 2^{\aleph_0}$ for h is a bijection. This completes the proof. ■

Corollary 9.4.1.1 *Theorem 9.4.1 holds, where G_C is replaced by a developmental paradigm.*

(4) *Let G_C be a denumerable general paradigm. There exists an intrinsic ultra-natural process, ${}^*\mathbf{S}$, such that objects described by members of G_C are produced by ${}^*\mathbf{S}$. During this production, numerous many pure NSP-objects as described by statements in ${}^*G_C - G_C$ are produced.*

.....

CHAPTER 9 REFERENCES

- 1 Evans, R., *The Atomic Nucleus*, McGraw-Hill, New York, 1955.
- 2 Mendelson, E., *Introduction to Mathematical Logic*, Ed. 2, D. Van Nostrand Co., New York, 1979.
- 3 Misner, Thorne and Wheeler, *Gravitation*, W. H. Freeman, San Francisco, 1973.
- 4 Patton C. and A. Wheeler, Is physics legislated by cosmogony? in *Quantum Gravity*, ed. Isham, Penrose and Sciama, Oxford University Press, Oxford, 1977, 538–605.

10.1 More About Ultrawords.

Previously, we slightly investigated the composition of an ultraword $w \in {}^*\mathbf{M} - \mathbf{d}$. Using the idea of the minimum informal language $P_0 \subset P$, where denumerable $B = \mathbf{d}$ and P is a propositional language, our interest now lies in completely determining the composition of ${}^*\mathbf{S}(\{w\})$. [Note: since our language is informal axiom (3) and (4) are redundant in that superfluous parentheses have been removed.] First, two defined sets.

$$(10.1.1) \quad A = \{x \mid x \in P_0 \text{ is an instance of an axiom for } S\}$$

$$(10.1.2) \quad C = \{x \mid x \in P_0 \text{ is a finite } (\geq 1) \text{ conjunction of members of } \mathbf{d}\}$$

Notice that it is also possible to refine the set C by considering C to be an ordered conjunction with respect to the ordering of the indexing set used to index members of $\mathbf{d}$. Further, as usual, we have that $A, C, \mathbf{d}$ are mutually disjoint.

Theorem 10.1.1 shows that the contents of the ultranatural events that are produced when the ultralogic ${}^*\mathbf{S}$ is applied to an ultraword w cannot be described in any standard human language. However, certain general properties can be described.

Theorem 10.1.1 *Let $w \in {}^*\mathbf{M}_{\mathbf{d}} - {}^*\mathbf{d}$ be an ultraword for infinite $\mathbf{d} \subset {}^*\mathbf{S}(\{w\})$. Then ${}^*\mathbf{S}(\{w\}) = {}^*\mathbf{A} \cup Q_1 \cup d'_1$, where for internal $*$ -finite d'_1 , $\mathbf{d} \subset d'_1 \subset {}^*\mathbf{d}$ and internal $Q_1 \subset {}^*\mathbf{C}$ is composed of $*$ -finite (≥ 1) conjunctions (i.e. $i(||| \text{and} |||)$) of distinct members of d'_1 and $w \in Q_1$. Further, each member of d'_1 and no other $*$ -propositions is used to form the $*$ -finite conjunctions in Q_1 , the only $*$ -propositions in ${}^*\mathbf{S}(\{w\})$ are those in w , and ${}^*\mathbf{A}$, Q_1 and d'_1 are mutually disjoint.*

Proof. The intent is to show that if $w \in \mathbf{M} - \mathbf{d}$, then $S(\{w\}) = A \cup Q \cup d'$, where $Q \subset C$, finite $d' \subset \mathbf{d}$ and Q is composed of finite (≥ 1) conjunctions of members of d' , each member of d' is used to form these conjunctions and no other propositions.

Let J be the set of propositional atoms in the composite w . (0) Then $J \subset S(\{w\})$. If K is the set of all propositional atoms in $S(\{w\})$, then $J \subset K$. Let $b \in K - J$. It is obvious that $b \notin S(\{w\})$ since otherwise $\{w, b\} \subset S_0(\{w\})$ but $\not\models_{S_0} w \rightarrow b$. Thus, $J = K$. Consequently, $J \subset S(\{w\})$, $J \subset \mathbf{d}$ and there does not exist an $F \in \mathbf{d} - J$, such that $F \in S(\{w\})$. (1) Let $J = d'$. The only propositional atoms in $S(\{w\})$ are those in w . Obviously $A \subset S(\{\emptyset\})$.

Assume the language P_0 is inductively defined from the set of atoms $\mathbf{d}$. Recall that for our axioms $\mathcal{X} = \mathcal{D} \rightarrow \mathcal{F}$, the strongest connective in $\mathcal{X}$ is $\rightarrow$. While in $\mathcal{D}$, or $\mathcal{F}$ when applicable, the strongest connective is $\wedge$. Since $\emptyset \subset \{w\}$, it follows that $S(\{w\}) = S(\emptyset) \cup S(\{w\})$. Let $b \in S(\emptyset)$. The only steps in the formal proof for b contain axioms or follows from modus ponens. Suppose that step $B_k = b$ is the first modus ponens step obtained from steps B_i, B_j , $i, j < k$, where $B_i = A \rightarrow b$, $B_j = A$. The strongest connective for each axiom is $\rightarrow$. However, since $A \rightarrow b$ is an axiom, the strongest connective in A is $\wedge$. This contradicts the requirement that A must also

be an axiom with strongest connect $\rightarrow$. Thus no modus ponens step can occur in a formal proof for b. Hence, (2) $A = S(\emptyset)$. (No modus ponens step can occur using two axioms.)

Let $B_k = b_1 \in P_0$ and suppose (a) that $b_1 = w$, or (b) $b_1 \neq w$ and is the first nonaxiom step that appears in a formal demonstration from the hypothesis w . Assume (b). Then all steps $B_i \in \{w\} \cup A$, $0 \leq i < k$. Then the only way that b_1 can be obtained is by means of modus ponens. However, all other steps, not including that which is w , are axioms. No modus ponens step can occur using two axioms. Thus one of the steps used for modus ponens must not be an axiom. The only nonaxiom that occurs prior to the step B_k is the step $B_m = w$. Hence, one of the steps required for B_k must be $B_m = w$. The other step must be an axiom of the form $w \rightarrow b_1$ and $b_1 \neq w$. Thus, from the definition of the axioms (3) b_1 is either a finite (≥ 1) conjunction of atoms in d' , or a single member of d' . Assume strong induction. Hence, for $n > 1$, statement (3) holds for all r , $1 \leq r \leq n$. A similar argument shows that (3) holds for the b_{n+1} nonaxiom step. Thus by induction, (3) holds for all nonaxiom steps.

Hence, there exists a $Q \subset C$ such that each member of Q is composed of finitely many (≥ 1) distinct members of d' and the set $G(Q)$ of all the proposition atoms that appear in any member of $Q = d' = J$ since $w \in Q$. Moreover, (4) $S(\{w\}) = A \cup d' \cup Q$ and (5) A , d' , Q are mutually disjoint.

$$\begin{aligned} \forall x(x \in \mathbf{M} - \mathbf{d} \rightarrow \exists y \exists z((y \in F(\mathbf{d})) \wedge (z \in \mathbf{C}) \wedge (S(\{x\}) = \\ \mathbf{A} \cup y \cup z) \wedge (\mathbf{A} \cap y = \emptyset) \wedge (\mathbf{A} \cap z = \emptyset) \wedge (x \in z) \wedge \\ (y \cap z = \emptyset) \wedge \mathbf{G}(z) = y)). \end{aligned} \quad (10.1.3)$$

holds in $\mathcal{M}$, hence also in $^*\mathcal{M}$. So, let w be an ultraword. Then there exists internal $Q_1 \subset ^*\mathbf{C}$, $w \in Q_1$ and * -finite $d'_1 \subset ^*\mathbf{d}$ such that $\mathbf{d} \subset ^*\mathbf{S}(\{w\}) = ^*\mathbf{A} \cup d'_1 \cup Q_1$; $^*\mathbf{A}$, d'_1 , Q_1 are mutually disjoint and $^*G(Q_1) = d'_1 = ^*\mathbf{J}$. Hence, $\mathbf{d} \subset d'_1$.

Now to analyze the objects in Q_1 . Let $\mathbf{d} = \{F_i \mid i \in \mathbb{N}\}$. Consider a bijection $h: \mathbb{N} \rightarrow \mathbf{d}$ defined by $h(n) = \mathbf{F}_n = [f]$, where $f \in T^0$ is the special member of $\mathbf{F}_n$ such that $f = \{(0, f(0))\}$, $f(0) = i(\mathbf{F}_n) = q_n \in i[\mathbf{d}]$. From the above analysis, (A) $[g] \in \mathbf{S}(\{w\}) - \mathbf{A} - \mathbf{d}$, ($w \in \mathbf{M}_\mathbf{d} - \mathbf{d}$), iff there exist $k, j \in \mathbb{N}$ such that $k < j$ and $f'_1 \in i[\mathbf{P}_0]^{2(j-k)}$ such that $[f'_1] = [g]$, and this leads to (B) that for each even $2p$, $0 \leq 2p \leq 2(j-k)$; $f'_1(2p) = q_{k+p} \in i[\mathbf{P}_0] \subset A_1$, $[(0, q_{k+p})] \in \mathbf{d}'$, all such q_{k+p} being distinct. For each odd $2p+1$ such that $0 \leq 2p+1 \leq 2(j-k)$, $f'_1(2p+1) = i(||| \text{and} |||)$. Also (C) $h(p) \in h[[k, j]]$ iff there exists an even $2p$ such that $0 \leq 2p \leq 2(j-k)$ and $f'_1(2p) = h(p) = q_{k+p} \in i[\mathbf{P}_0]$. [Note that 0 is considered to be an even number.]

By * -transfer of the above statements (A), (B) and (C), $[g] \in Q_1$ iff there exists some $j, k \in ^*\mathbb{N}$, $k < j$, and $f' \in ^*(i[\mathbf{P}_0])^{2(j-k)}$ such that $[f'] = [g]$ and $^*h[[k, j]] \subset ^*\mathbf{d}$. Moreover, each $^*h(r)$, $r \in [k, j]$ is a distinct member of $^*\mathbf{d}$. The conjunction “codes” for $i(||| \text{and} |||) \in A_1$ that are generated by each odd $2p+1$ are all the same and there are * -finitely many of them. Hence, Q_1 is the * -finite (≥ 1) conjunctions of distinct members of d'_1 , no other * -propositions are utilized and since $^*\mathbf{G}(Q_1) = d'_1$, all members of d'_1 are employed for these conjunctions. This completes the proof. ■

Corollary 10.1.1.1 *Let $w \in ^*\mathbf{M} - ^*\mathbf{d}$ be an ultraword for denumerable $\mathbf{d}$ such that $\mathbf{d} \subset ^*\mathbf{S}(\{w\})$. Then $^*\mathbf{S}(\{w\}) \cap \mathbf{P}_0 = \mathbf{A} \cup Q \cup \mathbf{d}$ and A , Q , $\mathbf{d}$ are mutually disjoint. The set Q is composed of finite ≥ 1 conjunctions of members of $\mathbf{d}$ and all of the members of $\mathbf{d}$ are employed to obtain these conjunctions.*

Proof. Recall that due to the finitary character of our standard objects $^\sigma \mathbf{A} = \mathbf{A} = ^*\mathbf{A} \cap \mathbf{P}_0$. In like manner, since $\mathbf{d} \subset d'_1$, $d'_1 \cap \mathbf{P}_0 = \mathbf{d}$. Now $\mathbf{P}_0 \cap Q_1$ are all of the standard members of Q_1 .

For each $k \in {}^*\mathbb{N}$, ${}^*h(k) = \mathbf{F}_k \in {}^*\mathbf{d}$ and conversely. Further, $\mathbf{F}_k \in \mathbf{d}$ iff $k \in \mathbb{N}$. Restricting $k, j \in \mathbb{N}$ in the above theorem yields standard finite ≥ 1 conjunctions of standard members of d'_1 ; hence, members of $\mathbf{d}$. Since ultraword $w \in Q_1$, we know that there exists some $\eta \in {}^*\mathbb{N} - \mathbb{N}$ and $f'_1 \in {}^*(i[P_0])^{2\eta}$, where f'_1 satisfies the * -transfer of the properties listed in the above theorem. Since finite conjunctions of standard members of d'_1 are * -finite conjunctions of members of d'_1 and $\mathbf{d} = \mathbf{d} \cap d'_1$, it follows that all possible finite conjunctions of members of $\mathbf{d}$ that are characterized by the function $f'_1 \in i[P_0]^{2(j-k)}$ are members of Q_1 for each such $j, k < \eta$. Also for such j, k the values of f'_1 are standard. On the other hand, any value of f'_1 is nonstandard iff it corresponds to a member of $d'_1 - \mathbf{d}$. Thus $Q_1 \cap \mathbf{P}_0 = \mathbf{Q}$ and this completes the proof. ■

If it is assumed that each member of $\mathbf{d}$ describes a Natural event (i.e. N-event) at times indicated by X_i , dropping the X_i may still yield a denumerable developmental paradigm without specifically generated symbols such as the “ i .” Noting that d'_1 is * -finite and internal leads to the conclusion that we can have little or no knowledge about the word-like construction of each member of $d'_1 - \mathbf{d}$. These pure nonstandard objects can be considered as describing pure NSP-world events, as will soon be demonstrated. Therefore, it is important to understand the following interpretation scheme, where descriptions are corresponded to events.

Standard or internal NSP-world events or sets of events are interpreted as directly or indirectly influencing N-world events. Certain external objects, such as the standard part operator, among others, are also interpreted as directly or indirectly influencing N-world events.

Notice that standard events can directly or indirectly affect standard events. In the micro-world, the term *indirect evidence* or verification is a different idea than indirect influences. You can have direct or indirect evidence of direct or indirect influences when considered within the N-world. An indirect influence occurs when there exists, or there is assumed to exist, a mediating “something” between two events. Of course, indirect evidence refers to behavior that can be observed by normally accepted human sensors as such behavior is assumed to be caused by unobserved events. However, the evidence for pure NSP-world events that directly or indirectly influence N-world events must be indirect evidence under the above interpretation.

In order to formally consider NSP-world events for the formation of objective standard reality, proceed as follows: let $\mathcal{O}$ be the subset of $\mathcal{W}$ that describes those Natural events that are used to obtain developmental or general paradigms and the like. Let $E_j \in \mathcal{O}$. Linguistically, assume that each E_j has the spacing symbol $|||$ immediately to the right. Thus within each T_i , there is a finite symbol string $F_i = E_i \in \mathcal{O}$ that can be joined by the juxtaposition (i.e. join) operator to other event descriptions. Assume that $\mathcal{W}_1$ is the set of nonempty symbol strings (with repetitions) formed (i.e. any finite permutation) from members of $\mathcal{O}$ by the join operator. These finite strings of symbols generate the basic elements for our partial sequences.

Obviously, $\mathcal{W}_1 \subset \mathcal{W}$. Consider $T'_i = \{XW_i \mid X \in \mathcal{W}_1\}$ and note that in many applications the time indicator W_i need not be of significance for a given E_j in some of the strings. Obviously, $T'_i \subset T_i$ for each i . For our isomorphism i onto A_1 , the following hold.

$$\begin{aligned}
 (10.1.4) \quad & \forall y(y \in \mathcal{E} \rightarrow (y \in \mathbf{T}'_i \leftrightarrow \exists x \exists f \exists w((\emptyset \neq w \in F(i[\mathcal{O}])) \wedge (x \in \mathbb{N}) \wedge \\
 & (f(0) = i[W_i]) \wedge (f \in P) \wedge \forall z((z \in \mathbb{N}) \wedge (0 < z \leq x) \rightarrow \\
 & f(z) \in w) \wedge (f \in y))))).
 \end{aligned}$$

$$\forall x(x \in \mathbb{N} \rightarrow \exists f \exists w((\emptyset \neq w \in F(i[\mathcal{O}])) \wedge (f \in P) \wedge \forall z(z \in \mathbb{N} \rightarrow (0 < z \leq x \leftrightarrow f(z) \in w))))). \quad (10.1.5)$$

$$\forall w(\emptyset \neq w \in F(i[\mathcal{O}]) \rightarrow \exists x \exists f \exists y((f \in P) \wedge (x \in \mathbb{N}) \wedge (y \in \mathbf{T}'_i) \wedge (f \in y) \wedge \forall z(z \in \mathbb{N} \rightarrow (0 < z \leq x \leftrightarrow f(z) \in w))))). \quad (10.1.6)$$

Since each finite segment of a developmental paradigm corresponds to a member of $\mathbf{T}'_i$, each nonfinite hyperfinite segment should correspond to a member of $*(\mathbf{T}'_i) - \mathbf{T}'_i$ and it should be certain individual segments of such members of $*(\mathbf{T}'_i) - \mathbf{T}'_i$ that correspond to the ultranatural events produced by an ultraword; UN-events that cannot be eliminated from an NSP-world developmental paradigms. [Note: for a scientific language, 10.1.4 - 10.1.6 and other such statements would correspond to an $\mathcal{W}'$ as generated by, at the least, a denumerable alphabet such as used in 9.2, 9.3.]

10.2 Laws and Rules.

One of the basic requirements of human mental activity is the ability to recognize the symbolic differences between finitely long strings of symbols as necessitated by our reading ability and to apply linguistic rules finitely many times. Gödel numberings specifically utilize such recognitions and the rules for the generation of recursive functions must be comprehended with respect to finitely many applications. Observe that Gödel number recognition is an “ordered” process while some fixed intuitive order is not necessary for the application of the rules that generate recursive functions.

In general, the simplest “rule” for ordered or unordered finite human choice, a rule that is assumed to be humanly comprehensible by finite recognition, is to simply *list* the results of our choice (assuming that they are symbolically representible in some fashion) as a partial finite sequence for ordered choice or as a finite set of finitely long symbol strings for an unordered choice. Hence, the end result for a finite choice can itself be considered as an algorithm “for that choice only.” The next application of such a *finite choice rule* would yield the exact same partial sequence or choice set. Another more general rule would be a statement which would say that you should “choose a specific number of objects” from a fixed set (of statements). Yet, a more general rule would be that you simply are required to “choose a finite set of all such objects,” where the term “finite” is intuitively known. Of course, there are numerous specifically described algorithms that will also yield finite choice sets.

From the symbol string viewpoint, there are trivial machine programmable algorithms that allow for the comparison of finitely long symbols with each member of a finite set of symbol strings B that will determine whether or not a specific symbol string is a member of B . These programs duplicate the results of human symbol recognition. As is well-known, there has not been an algorithm described that allows us to determine whether or not a given finite symbol string is a member of the set of all theorems of such theories as formal Peano Arithmetic. If one accepts Church’s Thesis, then no such algorithm will ever be described.

Define the general finite human choice relation on a set A as $H_0(A) = \{(A, x) \mid x \in F_0(A)\}$, where F_0 is the finite power set operator (including the empty set = no choice is made). Obviously, the inverse H_0^{-1} is a function from $F(A)$ onto $\{A\}$. There are choice operators that produce sets with a specific number of elements that can be easily defined. Let $F_1(A)$ be the set of all singleton subsets of A . The axioms of set theory state that such a set of singleton sets exists. Define $H_1(A) = \{(A, x) \mid x \in F_1(A)\}$, etc. Considering such functions as defined on sets X that are members of a superstructure, then these relations are subsets of $\mathcal{P}(X) \times \mathcal{P}(X)$ and as such are also members of the superstructure.

The next discussion shows that the IUN-selection processes ${}^*\mathbf{C}_i$, $i \geq 0$ exist as formal objects within the NSP-world.

Let $\mathbf{A} = \mathbf{P}_0$. Observe that ${}^\sigma\mathbf{H}_0(\mathbf{A}) = \{({}^*\mathbf{A}, x) \mid x \in F_0(\mathbf{A})\}$ and ${}^*\mathbf{H}_1(\mathbf{A}) = \{({}^*\mathbf{A}, x) \mid x \in {}^*(F_1(\mathbf{A}))\}$ ($i \geq 0$). Now ${}^*(F_0(\mathbf{A})) = {}^*F_0({}^*\mathbf{A})$ is the set of all * -finite subsets of ${}^*\mathbf{A}$. On the other hand, for the $i > 0$ cardinal subsets, ${}^*(F_i(\mathbf{A})) = F_i({}^*\mathbf{A})$ for each $i \geq 1$. With respect to an ultraword w that generates the general and developmental paradigms, we know that $w \in {}^*\mathbf{P}_0 - \mathbf{P}_0$ and that $({}^*\mathbf{P}_0, \{x\}) \in {}^*\mathbf{H}_1(\mathbf{P}_0)$. The actual finite choice operators are characterized by the set-theoretic second projector operator P_2 as it is defined on $H_i(\mathbf{A})$. This operator embedded by the injection θ is the same as P_2 as it is defined on $\mathbf{H}_i(\mathbf{A})$. Thus, when $h = (\mathbf{A}, x) \in \mathbf{H}_i(\mathbf{A})$, then we can define $x = P_2(h) = C_i(h) = \mathbf{C}_i(h)$. The maps C_i and $\mathbf{C}_i$, formally defined below, are the specific finite choice operators. For consistency, we let C_i and $\mathbf{C}_i$ denote the appropriate finite choice operators for $H_i(\mathbf{A})$ and $\mathbf{H}_i(\mathbf{A})$, respectively.

Since the *P_2 defined on say $\mathbf{H}_i(\mathbf{A})$ is the same as the set-theoretic second projection operator P_2 , it would be possible to denote ${}^*\mathbf{C}_i$ as $\mathbf{C}_i$ on internal objects. For consistency, the notation ${}^*\mathbf{C}_i$ for these special finite choice operators is retained. Formally, let $\mathbf{C}_i: \mathbf{H}_i(\mathbf{A}) \rightarrow F_i(\mathbf{A})$. Observe that ${}^\sigma\mathbf{C}_i = \{({}^*(a, b) \mid (a, b) \in \mathbf{C}_i\} = \{({}^*((\mathbf{A}, b), b) \mid b \in F_i(\mathbf{A}))\} \subset {}^*\mathbf{C}_i$; and, for $b \in F_i(\mathbf{A})$, $\mathbf{C}_i((\mathbf{A}, b)) = b$ implies that ${}^\sigma(\mathbf{C}_i((\mathbf{A}, b))) = \{{}^*a \mid a \in b\} = b$ from the construction of $\mathcal{E}$. Thus in contradistinction to the consequence operator, for each $({}^*\mathbf{A}, b) \in {}^*\mathbf{H}_i$, the image $({}^\sigma\mathbf{C}_i)(({}^*\mathbf{A}, b)) = {}^\sigma(\mathbf{C}_i((\mathbf{A}, b))) = ({}^*\mathbf{C}_i)(({}^*\mathbf{A}, b)) = b$. Consequently, the set map ${}^\sigma\mathbf{C}_i: {}^\sigma\mathbf{H}_i \rightarrow F_i(\mathbf{A}) = {}^\sigma(F_i(\mathbf{A}))$ and ${}^*\mathbf{C}_i \mid {}^\sigma\mathbf{H}_i = {}^\sigma\mathbf{C}_i$. Finally, it is not difficult to extend these finite choice results to general internal sets.

In the proofs of such theorems as 7.2.1, finite and other choice sets are selected due to their set-theoretic existence. The finite choice operators C_i are not specifically applied since these operators are only intended as a mathematical model for apparently effective human processes — procedures that generate acceptable algorithms. As is well-known, there are other describable rules that also lead to finite or infinite collections of statements. Of course, with respect to a Gödel encoding i for the set of all words $\mathcal{W}$ the finite choice of readable sentences in $\mathcal{E}$ is one-to-one and effectively related to a finite and, hence, recursive subset of $\mathbb{N}$.

From this discussion, the descriptions of the finite choice operators would determine a subset of the set of all algorithms (“rules” written in the language $\mathcal{W}$) that allow for the selection of readable sentences. Notice that before algorithms are applied there may be yet another set of readable sentences that yields conditions that must exist prior to an application of such an algorithm and that these application rules can be modeled by members of $\mathcal{E}$.

In order to be as unbiased as possible, it has been required for N-world applications that the set of all frozen segments be infinite. Thus, within the proof of Theorem 7.2.1, every N-world developmental, as well as a general paradigm, is a proper subset of a * -finite NSP-world paradigm, and the * -finite paradigm is obtained by application of the * -finite choice operator ${}^*\mathbf{C}_0$. As has been shown, such * -finite paradigms contain pure unreadable (subtle) sentences that may be interpreted for developmental paradigms as pure refined NSP-world behavior and for general paradigms as specific pure NSP-world ultranatural events or objects.

Letting Γ correspond to the formal theory of Peano Arithmetic, then assuming Church’s Thesis, there would not exist a N-world algorithm (in any human language) that allows for the determination of whether or not a statement F in the formal language used to express Γ is a member of Γ . By application of the * -finite choice operator ${}^*\mathbf{C}_0$, however, there does exist a * -finite Γ' such that

$\sigma\Gamma = \Gamma \subset \Gamma'$ and, hence, within the NSP-world a “rule” that allows the determination of whether or not $F \in \Gamma'$. If such internal processes mirror the only allowable procedures in the NSP-world for such a “rule,” then it might be argued that we do not have an effective NSP-world process that determines whether or not F is a member of Γ for Γ is external.

As previously alluded to at the beginning of this section, when a Gödel encoding i is utilized with the N-world, the injection i is not a surjection. When such Gödel encodings are studied, it is usually *assumed*, without any further discussion, that there is some human mental process that allows us to recognize that one natural number representation (whether in prime factored form or not) is or is not distinct from another such representation. It is not an unreasonable assumption to assume that the same *effective* (but external) process exists within the NSP-world. Thus within the NSP-world there is a “process” that determines whether or not an object is a member of ${}^*\mathbb{N} - \mathbb{N} = \mathbb{N}_\infty$ or $\mathbb{N}$. Indeed, from the ultraproduct construction of our nonstandard model, a few differences can be detected by the human mathematician. Consequently, this assumed NSP-world effective process would allow a determination of whether or not $\mathbf{F} = [f_m]$ is a member of Γ by recalling that $f_m \in P_m$ signifies that $[f_m] \in {}^*\Gamma - \Gamma$ implies $m \in \mathbb{N}_\infty \simeq {}^*(i[\mathcal{W}]) - i[\mathcal{W}] = {}^*A_1 - A_1$.

The above NSP-world recognition process is equivalent, as defined in Theorem 7.2.1, to various applications of a single (external) set-theoretic intersection. Therefore, there are internal processes, such as ${}^*\mathbf{C}_0$, that yield pure NSP-world developmental paradigms and a second (external) but acceptable NSP-world effective process that produces specific N-world objects. Relative to our modeling procedures, it can be concluded that both of these processes are intrinsic ultranatural processes.

With respect to Theorem 10.1.1, the NSP-world developmental or general paradigm generated by an ultraword is * -finite and, hence, specifically NSP-world obtainable prior to application of ${}^*\mathbf{S}$ through application of ${}^*\mathbf{C}_0$ to ${}^*\mathbf{d}$. However, this composition can be reversed. The NSP-world (IUN) process ${}^*\mathbf{C}_1$ can be applied to the appropriate ${}^*\mathbf{M}$ type set and an appropriate ultraword $w \in {}^*\mathbf{M}$ obtained. Composing ${}^*\mathbf{C}_1$ with ${}^*\mathbf{S}$ would yield d'_1 in a slightly less conspicuous manner. Obviously, different ultrawords generate different standard and nonstandard developmental or general paradigms.

To complete the actual mental-type processes that lead to the proper ordered event sequences, the above discussion for the finite choice operators is extended to the human mental ability of ordering a finite set in terms of rational number subscripts. New choice operators are defined that model not just the selection of a specific set of elements that is of a fixed finite cardinality but also choosing the elements in the required rational number ordering. The ultrawords w that exist are * -finite in length. By application of the inverses of the f and τ functions of section 7.1, where they may be considered as extended standard functions *f and ${}^*\tau$, there would be from analysis of extended theorem 7.3.2 a hyperfinite set composed of standard or nonstandard frozen segments contained in an ultraword. Further, in theorem 7.3.2, the chosen function f does not specifically differentiate each standard or nonstandard frozen segment with respect to its “time” stamp subscript. There does exist, however, another function in the * -equivalence class $[g] = w$ that will make this differentiation. It should not be difficult to establish that after application of the ultralogic ${}^*\mathbf{S}$, there is applied an appropriate mental-like hyperfinite ordered choice operator (an IUN-selection process) and that this would yield that various types of event sequences. Please note that each event sequence has a beginning point of observation. This point of observation need not indicate the actual moment when a specific Natural-system began its development.

In the following discussion, we analyze how ultranatural laws aid in the production of Natural and UN-event sequences, and develop an external unification of all physical theories.

Various subdevelopmental (or subgeneral) paradigms d_i are obtained by considering the actual descriptive content (i.e. events) of specific theories Γ_i that are deduced from hypotheses η_i , usually, by finitary consequence operators S_i (the inner logics) that are compatible with S . In this case, $d_i \subset S_i(\eta_i)$. It is also possible to include within $\{d_i\}$ and $\{\eta_i\}$ the assumed descriptive chaotic behavior that seems to have no apparent set of hypotheses except for that particular developmental paradigm itself and no apparent deductive process except for the identity consequence operator. In this way, such scientific nontheories can still be considered as a formal theory produced by a finitary consequence operator applied to an hypothesis. Many of these hypotheses η_i contain the so-called natural laws (or first-principles) peculiar to the formal theories Γ_i and the theories language, where it is assume that such languages are at least closed under the informal conjunction and conditional.

Consider each η_i to be a general paradigm. For the appropriate M type set constructed from the denumerable set $B = \{\bigcup\{d_i\} \cup (\bigcup\{\eta_i\})$, redefine M_B to be the smallest subset of P_0 containing B and closed under finite (≥ 0) conjunction. (The usual type of inductively defined M_B .) Then there exist ultrawords $w_i \in {}^*M_B - {}^*B$ such that $\eta_i \subset {}^*S(\{w_i\})$ (where due to parameters usually *ultranatural laws* exist in ${}^*S(\{w_i\}) - \eta_i$ and $d_i \subset {}^*S(\{w_i\})$. Using methods such as those in Theorem 7.3.4, it follows that there exists some $w'' \in {}^*M_B - M_B$ such that $w_i \in {}^*S(\{w''\})$ and, consequently, $\eta_i \cup d_i \subset {}^*S(\{w''\})$. Linguistically, it is hard to describe the ultraword w'' . Such a w'' might be called an *ultimate ultranatural hypothesis* or *the ultimate building plain*.

Remark. It is not required that the so-called Natural laws that appear in some of the η_i be either cosmic time or universally applicable. They could refer only to local first-principles. It is not assumed that those first-principles that display themselves in our local environment are universally space-time valid.

Since the consequence operator S is compatible with each S_i , it is useful to proceed in the following manner. First, apply the IUN-process *S to $\{w''\}$. Then $d_i \cup \eta_i \subset {}^*S(\{w''\})$. It now follows that $d_i \cup \eta_i \subset {}^*S(\{w_i\}) \subset {}^*S_i({}^*S(\{w_i\})) \subset {}^*S_i({}^*S_i(\{w_i\})) = {}^*S_i(\{w_i\})$. Observe that for each $a \in \Gamma_i$ there exists some finite $F_i \subset \eta_i$ such that $a \in S_i(F_i)$. However, $F_i \subset \eta_i$ for each member of $F(\eta_i)$ implies that $a \in {}^*S_i(F_i) \subset {}^*S_i({}^*S(\{w_i\}))$. Consequently, $\Gamma_i \subset {}^*S_i({}^*S(\{w_i\}))$. The ultimate ultraword suffices for the descriptive content and inner logics associated with each theory Γ_i .

We now make the following observations relative to “rules” and deductive logic. It has been said that science is a combination of empirical data, induction and deduction, and that you can have the first two without the last. That this belief is totally false should be self-evident since the philosophy of science requires its own general rules for observation, induction, data collection, proper experimentation and the like. All of these general rules require logical deduction for their application to specific cases — the metalogic. Further, there are specific rules for linguistics that also must be properly applied prior to scientific communication. Indeed, we cannot even open the laboratory door — or at least describe the process — without application of deductive logic. The concept of deductive logic as being the patterns our “minds” follow and its use exterior to the inner logic of some theory should not be dismissed for even the (assumed?) mental methods of human choice that occur prior to communicating various scientific statements and descriptions.

Finally, with respect to the hypothesis rule in [9], it might be argued that we can easily analyze

the specific composition of all significant ultrawords, as has been previously done, and the composition of the nonstandard extension of the general paradigm. Using this assumed analysis and an additional alphabet, one *might* obtain specific information about pure NSP-world ultranatural laws or refined behavior. Such an argument would seem to invalidate the cautious hypothesis rule and lead to appropriate speculation. However, such an argument would itself be invalid.

Let $\mathcal{W}_1$ be an infinite set of meaningful readable sentences for some description and assume that $\mathcal{W}_1$ does not contain any infinite subset of readable sentences each one of which contains a mathematically interpreted entry such as a real number or the like. Since $\mathcal{W}_1 \subset \mathcal{W}$ and the totality $\mathbf{T}_i = \{XW_i \mid X \in \mathcal{W}\}$ is denumerable, the subtotality $\mathbf{T}'_i = \{XW_i \mid X \in \mathcal{W}_1\}$ is also denumerable. Hence, the external cardinality of ${}^*\mathbf{T}'_i \geq 2^{|\mathcal{M}|}$.

Consider the following sentence

$$\begin{aligned} \forall z(z \in i[\mathcal{W}_1] \rightarrow \exists y \exists x((y \in A_1^{[0,1]}) \wedge (x \in \mathbf{T}'_i) \wedge (y \in x) \wedge \\ ((0, i(W_i)) \in y) \wedge ((1, z) \in y))). \end{aligned} \quad (10.2.1)$$

By *-transfer and letting “z” be an element in ${}^*(i[\mathcal{W}_1]) - i[\mathcal{W}_1]$ it follows that we can have little knowledge about the remaining and what must be unreadable portions that take the “X” position. If one assumes that members of $\mathcal{W}_1$ are possible descriptions for possible NSP-world behavior at the time t_i , then it may be assumed that at the time t_i the members of ${}^*\mathbf{T}'_i - \mathbf{T}_i$ describe NSP-world behavior at NSP-world (and N-world) time t_i . Now as i varies over ${}^*\mathbb{N}$, pure nonstandard subdevelopmental paradigms (with or without the time index statement W_i) exist with members in ${}^*\mathcal{T}$ and may be considered as descriptions for time refined NSP-world behavior, especially for a NSP-world time index $i \in \mathbb{N}_\infty$.

CHAPTER 10 REFERENCES

- 1 Beltrametti, E. G., Enrico, G. and G. Cassinelli, *The Logic of Quantum Mechanics*, Encyclopedia of Mathematics and Its Applications, Vol. 15, Addison-Wesley, Reading, 1981.
- 2 d’Espagnat, B., The quantum theory and realism, *Scientific America*, 241(5)(1979), 177.
- 3 *Ibid.*
- 4 *Ibid.*, 181.
- 5 *Ibid.*,
- 6 *Ibid.*, 180.
- 7 Feinberg, G., Possibility of faster-than-light particles, *Physical Review*, 159(5)(1976), 1089—1105.
- 8 Hanson, W. C., The isomorphism property in nonstandard analysis and its use in the theory of Banach Spaces, *J. of Symbolic Logic*, 39(4)(1974), 717—731.
- 9 Herrmann, R. A., D-world evidence, *C.R.S. Quarterly*, 23(2)(1986), 47—54.
- 10 Herrmann, R. A., The Q-topology, Whyburn type filters and the cluster set map, *Proceedings Amer. Math. Soc.*, 59(1)(1975), 161—166.
- 11 Kleene, S. C., *Introduction to Metamathematics*, D. Van Nostrand Co., Princeton, 1950.
- 12 Prokhovnik, S. J., *The Logic of Special Relativity*, Cambridge University Press, Cambridge, 1967.
- 13 Stroyan, K. D. and W. A. J. Luxemburg, *Introduction to the Theory of Infinitesimals*, Academic Press, New York, 1976.
- 14 Tarski, A., *Logic, Semantics, Metamathematics*, Clarendon Press, Oxford, 1969.

15 Thurber, J. K. and J. Katz, Applications of fractional powers of delta functions, *Victoria Symposium on Nonstandard Analysis*, Springer-Verlag, New York, 1974.

16 Zakon, E., Remarks on the nonstandard real axis, *Applications of Model Theory to Algebra, Analysis and Probability*, Holt, Rinehart and Winston, New York, 1969.

17 Note that the NSP-world model is not a local hidden variable theory.

In the next section, it is shown how the mathematical structure predicts the possible existence of the ultrasubparticle and how, by the very simple process of ultrafinite combinations, all of the Natural universe assumed fundamental entities can be produced. Also, this allows for the construction of infinitely many distinct collections of fundamental entities for distinct universes. Further, it allows the virtual particle concept to be replaced by ultrafinite combinations.

11.1 Subparticles.

What is a subparticle? Or, what is an infant? Or, better still, what is a thing? I first used the name infant for these strange objects. I then coined the term subparticle. It would have been better to have simply called them “things.” As stated in [9], these objects are not to be described in terms of any geometric configuration. These multifaceted things, these subparticles, are not to be construed as either particles nor waves nor quanta nor anything that can be represented by some fixed imagery. Subparticles are to be viewed only operationally. Subparticles are only to be considered as represented by a *-finite sequence $\{a_i\}_{i=1}^n$, $n \in {}^*\mathbb{N}$, of hyperreal numbers. Indeed, the idea of the n -tuple $(a_1, a_2, \dots, a_i, \dots)$ notation is useful and we assume that n is a fixed member of $\mathbb{N}_\infty$. The language of coordinates for this notation is used, where the i 'th coordinate means the i 'th value of the sequence. Obviously, 0 is not a domain member for our sequential representation.

The first coordinate a_1 is a “naming” coordinate. The remaining coordinates are used to represent various real numbers, complex numbers, vectors, and the like physical qualities needed for different physical theories. For example, $a_2 = 1$ might be a counting coordinate. Then a_i , $3 \leq i \leq 6$ are hyperreal numbers that represent NSP-world coordinate locations of the subparticle named by a_1 — a_7 , a_8 represent the positive or negative charges that can be assigned to every subparticle — a_9 , a_{11} , a_{13} hyperreal representations for the inertial, gravitational and intrinsic (rest) mass. For vector quantities, continue this coordinate assignment and assign specific coordinate locations for the vector components. So as not to be biased, include as other coordinates hyperreal measures for qualities such as energy, apparent momentum, and all other physical qualities required within theories that must be combined in order to produce a reasonable description for N-world behavior. For the same reason, we do not assume that such N-world properties as the uncertainty principle hold for the NSP-world.

It is purposely assumed that the qualities represented by the coordinate a_i , $i \geq 3$ are not inner-related, in their basic construction, by any mathematical relation since it is such inner-relations that are assumed to mirror the N-world laws that govern the development of not only our present universe but previous as well as future developmental alterations. The same remarks apply to any possible and distinctly different universes that may or not occur. Thus, for these reasons, we view the subparticle as being totally characterized by such a sequence $\{a_i\}$ and always proceed cautiously when any attempt is made to describe all but the most general subparticle behavior. Why have we chosen to presuppose that subparticles are characterized by sequences, where the coordinates are hyperreal numbers?

For chapters 11, 12 assume EGS. Let r be a positive real number. The number r can be represented by a decimal number, where for uniqueness, the repeated 9s case is used for all terminating decimals. From this, it is seen that there is a sequence S_i of natural numbers such that $S_i/10^i \rightarrow r$. Consequently, for any $\omega \in \mathbb{N}_\infty = {}^*\mathbb{N} - \mathbb{N}$, it follows that $\pm {}^*S_\omega/10^\omega \in \mu(\pm r)$, where ${}^*S_\omega \in {}^*\mathbb{N}$ and $\mu(\pm r)$ is the monad about r . In [9], it is assumed that each coordinate a_i , $i \geq 3$ is characterized by the numerical quantity $\pm 10^{-\omega}$, $\omega \in \mathbb{N}_\infty$. Obviously, we need not confine ourselves to the number $10^{-\omega}$. Indeed, the next theorem has interesting applications relative to such a selection.

Theorem 11.1.1 *Let $\omega \in \mathbb{N}_\infty$. Then for each $r \in \mathbb{R}$ there exists an $x \in \{m/\omega \mid (m \in {}^*\mathbb{Z}) \wedge (|m| < \omega^2)\}$ such that $x \approx r$ (i.e. $x \in \mu(r)$).*

Proof. Let $r \in \mathbb{R}$. Then there exists some integer $n \in \mathbb{Z}$ such that $n \leq r < n + 1$. Partition $[n, n + 1]$ as follows: let $0 \neq m \in \mathbb{N}$ and consider $n \leq n + 1/m \leq \dots \leq n + (m - 1)/m \leq n + 1$. Then there exists a unique $a \in \{0, 1, \dots, m - 1\}$ such that $r \in [n + a/m, n + (a + 1)/m)$. For $m = 1$, let $S_1 = n = f_1/1$. For $m \geq 2$, if $a \in \{0, 1, \dots, m - 2\}$, then let $S_m = n + (a + 1)/m = (nm + a + 1)/m = f_m/m$; if $a = m - 1$, then let $S_m = n + (m - 1)/m = (nm + m - 1)/m = f_m/m$. This yields two sequences $S: \mathbb{N} \rightarrow \mathbb{Q}$ and $f: \mathbb{N} \rightarrow \mathbb{Z}$. It follows easily that $S_i \rightarrow r$. Hence, for each $\omega \in \mathbb{N}_\infty$, ${}^*S_\omega = {}^*f_\omega/\omega \approx r$ and ${}^*f_\omega \in {}^*\mathbb{Z}$. Observe that ${}^*f_\omega/\omega$ is a finite (i.e. limited) number. Hence, $|{}^*f_\omega/\omega| < \omega$ entails that $|{}^*f_\omega| < \omega^2$. Therefore, ${}^*f_\omega/\omega \in \{m/\omega \mid (m \in {}^*\mathbb{Z}) \wedge (|m| < \omega^2)\}$. ■

Corollary 11.1.1.1 *Let $\omega \in \mathbb{N}_\infty$. Then there exists a sequence $f: \mathbb{N} \rightarrow \mathbb{Z}$ such that ${}^*f_\omega/{}^*m_\omega \in \mu(r)$ for each $r \in \mathbb{R}$. Also, there exists a sequence $g: \mathbb{N} \rightarrow \mathbb{N}$ such that for each real $r \geq 0$, $(\pm {}^*g_\omega/\omega) \approx \pm r$.*

For the *ultrasubparticle*, each coordinate $a_i = 1/10^\omega$ $i \geq 3$ and odd, $a_i = -1/10^\omega$ $i \geq 4$ and even, $\omega \in \mathbb{N}_\infty$. From the above theorem, the choice of $10^{-\omega}$ as the basic numerical quantity is for convenience only and is not unique except in its infinitesimal character. Of course, the sequences chosen to represent the ultrasubparticle are pure internal objects and as such are considered to directly or indirectly affect the N-world. Why might the *-finite “length” of such subparticles (here is where we have replaced the NSP-world entity by its corresponding sequence) be of significance?

First, since our N-world languages are formed from a finite set of alphabets, it is not unreasonable to assume that NSP-world “languages” are composed from a *-finite set of alphabets. Indeed, since it should not be presupposed that there is an upper limit to the N-world alphabets, it would follow that the basic NSP-world set of alphabets is an infinite *-finite set. Although the interpretation method that has been chosen does not require such a restriction to be placed upon NSP-world alphabets, it is useful, for consistency, to assume that descriptions for substratum processes that affect, in either a directly or indirectly detectable manner, N-world events be so restricted. For the external NSP-world viewpoint, all such infinite *-finite objects have a very significant common property. [Note relative to the basic book “The Theory of Ultralogics.” In what follows $\mathcal{M}_1$ is the standard superstructure constructed on page 76 and not the object defined on page 57.]

Theorem 11.1.2 *All infinite *-finite members of our (ultralimit) model ${}^*\mathcal{M}_1$ have the same external cardinality which is $\geq |\mathcal{M}_1|$.*

Proof. Hanson [8] and Zakon [16] have done all of the difficult work for this result to hold. First, one of the results shown by Henson is that all infinite *-finite members of our ultralimit model have the same external cardinality. Since our model is a comprehensive enlargement, Zakon’s theorem 3.8 in [16] applies. Zakon shows that there exists a *-finite set, A , such that $|A| \geq |\mathcal{M}_1| = |\mathcal{R}|$. Since

A is infinite, Hanson's result now implies that all infinite $*$ -finite members of our model satisfy this inequality.■

For an extended infinite standard set $*A$ it is well-known that $|*A| \geq 2^{|\mathcal{M}_1|}$. One may use these various results and establish easily that there exist more than enough subparticles to obtain all of the cardinality statements relative to the three substratum levels that appear in [9] even if we assume that there are a continuum of finitely many subparticle qualities that are needed to create all of the N-world.

Consider the following infinite set of statements expressed in an extended alphabet.

$$(11.1.1) \quad \begin{aligned} G_A = \{ & \text{An} || \text{elementary} || \text{particle} || k'(i', j') || \text{with} || \\ & \text{total} || \text{energy} || c'+1/(n'). \mid ((i, j, n) \in \\ & \mathbb{N}^+ \times \mathbb{N}^+ \times \mathbb{N}^+) \wedge (1 \leq k \leq m) \}, \end{aligned}$$

where $\mathbb{N}^+$ is the set of all nonzero natural numbers and $m \in \mathbb{N}^+$. Applying the same procedure that appears in the proof of Theorem 9.3.1 and with a NSP-world alphabet, we obtain

$$(11.1.2) \quad \begin{aligned} G'_A = \{ & \text{An} || \text{elementary} || \text{particle} || k'(i', j') || \text{with} || \\ & \text{total} || \text{energy} || c'+1/(n'). \mid ((i, j, n) \in \\ & *\mathbb{N}^+ \times *\mathbb{N}^+ \times *\mathbb{N}^+) \wedge (1 \leq k \leq m) \}, \end{aligned}$$

Assume that there is at least one type of elementary particle with the properties stated in the set G_A . It will be shown in the next section that within the NSP-world there may be simple properties that lead to N-world energy being a manifestation of mass. For $c = 0$, we have another internal set of descriptions that forms a subset of G'_A .

$$(11.1.3) \quad \begin{aligned} & \{ \text{An} || \text{elementary} || \text{particle} || k'(i', j') || \text{with} || \\ & \text{total} || \text{energy} || c'+1/(10^{\gamma'}). \mid ((i, j, \gamma) \in \\ & *\mathbb{N}^+ \times *\mathbb{N}^+ \times *\mathbb{N}^+) \wedge (1 \leq k \leq m) \}, \end{aligned}$$

For our purposes, (11.1.3) leads immediately to the not ad hoc concept of subparticles with infinitesimal proper mass. As will be shown, such infinitesimal proper mass can be assumed to characterize any possible zero proper mass N-world entity. The set G'_A has meaning if there exists at least one natural entity that can possess the energy expressed by G_A , where this energy is measured in some private unit of measure.

Human beings combine together finitely many sentences to produce comprehensible descriptions. Moreover, all N-world human construction requires the composition of objectively real N-world objects. We model the idea of *finite composition* or *finite combination* by an N-world process. This produces a corresponding NSP-world intrinsic ultranatural process *ultrafinite composition* or *ultrafinite combination* that can either directly or indirectly affect the N-world, where its effect is indirectly inferred.

Let the index j vary over a hyperfinite interval and fix the other indices. Then the set of sentences

$$(11.1.4) \quad \begin{aligned} G''_A = \{ & \text{An} || \text{elementary} || \text{particle} || k'(i', j') || \text{with} || \\ & \text{total} || \text{energy} || c'+1/(n'). \mid (j \in *\mathbb{N}^+) \\ & \wedge (1 \leq j \leq \lambda) \}, \end{aligned}$$

where $\lambda \in {}^*\mathbb{N}^+$, $3 \leq i \in {}^*\mathbb{N}$, $n \in \mathbb{N}_\infty$ and $1 \leq k \leq m$, forms an internal linguistic object that can be assumed to describe a hyperfinite collection of ultranatural entities. Each member of G''_A has the i 'th coordinate that measures the proper mass and is infinitesimal (with respect to NSP-world private units of measure). In the N-world, finite combinations yield an event. Thus, with respect to such sets as G''_A , one can say that there are such N-world events iff there are ultrafinite combinations of NSP-world entities. And such ultrafinite combinations yield a NSP-world event that is an ultranatural entity.

Associated with such ultrafinite combinations for the entities described in G''_A there is a very significant procedure that yields the i 'th coordinate value for the entity obtained by such ultrafinite combinations. Such entities are called *intermediate subparticles*. Let $m_0 \geq 0$ be the N-world proper mass for an assumed elementary particle denoted by k' . If $m_0 = 0$, then let $\lambda = 1$. Otherwise, from Theorem 11.1.1, we know that there is a $\lambda \in {}^*\mathbb{N}$ such that $\lambda/(10^\omega) \in \mu(m_0)$, where $\omega \in \mathbb{N}_\infty$ and since $m_0 \neq 0$, $\lambda \in \mathbb{N}_\infty$. Consequently, for $b_n = 10^{-\omega}$, the *-finite sum

$$\sum_{n=1}^{\lambda} b_n = \sum_{n=1}^{\lambda} \frac{1}{10^\omega} = \frac{\lambda}{10^\omega} \quad (11.1.5)$$

has the property that $\mathbf{st}(\sum_{n=1}^{\lambda} 1/(10^\omega)) = m_0$. (Note the special summation notation for a constant summand.) The standard part operator $\mathbf{st}$ is an important external operator that is a continuous [11] NSP-world process that yields N-world effects. The appropriate interpretation is that

ultrafinite combinations of ultrasubparticles yield an intermediate subparticle that, after application of the standard part operator, has the same effect as an elementary particle with proper mass m_0 .

An additional relevant idea deals with the interpretation that the *-finite set G''_A exists at, say, nonstandard time, and that such a set is manifested at standard time when the operator $\mathbf{st}$ is applied. The standard part operator is one of those external operators that can be indirectly detected by the presence of elementary particles with proper mass m_0 .

The above discussion of the creation of intermediate subparticles yields a possible manner in which ultrasubparticles are combined within the NSP-world to yield appropriate energy or mass coordinates for the multifaceted subparticles. But is there an indication that all standard world physical qualities that are denoted by qualitative measures begin as infinitesimals?

Consider the infinitesimal methods used to obtain such things as the charge on a sphere, charge density and the like. In all such cases, it is assumed that charge can be infinitesimalized. In 1972, it was shown how a classical theory for the electron, when infinitesimalized, leads to the point charge concept of quantum field theory and then how the *-finite many body problem produced the quasi-particle. [15] Although this method is not the same as the more general and less ad hoc subparticle approach, it does present a procedure that leads to an infinitesimal charge density and then, in a very ad hoc manner, it is assumed that there are objects that when *-finitely combined together entail a real charge and charge density. Further, it is the highly successful use of the modeling methods of infinitesimal calculus over hundreds of years that has lead to our additional presumption that all coordinates of the basic sequential subparticle representation are a $\pm$ fixed infinitesimal.

In order to retain the general independence of the coordinate representation, *independent *-finite coordinate summation* is allowed, recalling that such objects are to be utilized to construct

many possible universes. [This is the same idea as *-finitely repeated simple affine or linear transformations.] Thus, distinct from coordinatewise addition, *-finitely many such sequences can be added together by means of a fixed coordinate operator in the following sense. Let $\{a_i\}$ represent an ultrasubparticle. Fix the coordinate j , then the sequence $\{c_i\}$, $c_i = a_i$, $i \neq j$ and $c_j = 2a_j$ forms an intermediate subparticle. As will be shown, it is only after the formation of such intermediate subparticles that the customary coordinatewise addition is allowed and this yields, after the standard part operator is applied, representations for elementary particles. Hence, from our previous example, we have that ultrafinite combinations of ultrasubparticles yield subparticles with “proper mass” $\lambda/(10^\omega) \approx m_0$ while all other coordinates remain as $\pm 10^\omega$. This physical-like process is not a speculative ad hoc construct, but, rather, it is modeled after what occurs in our observable natural world. Intuitively, this type of summation is modeled after the process of inserting finitely many pieces of information (mail) into a single “postal box,” where these boxes are found in rectangular arrays in post offices throughout the world.

Now other ultrasubparticles are ultrafinitely combined and yield for a specific coordinate the $\pm$ unite charge or, if quarks exist, other N-world charges, while all other coordinates remain fixed as $\pm 1/(10^\omega)$, etc. Rationally, how can one conceive of a combination of these intermediate subparticles, a combination that will produce entities that can be characterized in a standard particle or wave language?

Recall that a finite summation is a *-finite summation within the NSP-world. Therefore, a finite combination of intermediate subparticles is an allowed internal process. [Note that external processes are always allowed but with respect to our interpretation procedures we always have direct or indirect knowledge relative to application of internal processes. Only for very special and reasonable external processes do we have direct or indirect knowledge that they have been applied.] Let $\gamma_i \in \mu(0)$, $i = 1, \dots, n$. Then $\gamma_1 + \dots + \gamma_n \in \mu(0)$. The final stage in subparticle formation for our universe — the final stage in particle or wave substratum formation — would be *finite coordinatewise* summation of finitely many intermediate subparticles. This presupposes that the N-world environment is characterized by but finitely many qualities that can be numerically characterized. This produces the following type of coordinate representation for a specific coordinate j after n summations with n other intermediate subparticles that have only infinitesimals in the j coordinate position.

$$(11.1.6) \quad \sum_{i=1}^{\lambda} (1/(10^\omega)) + \sum_{i=1}^n \gamma_i.$$

Assuming λ is one of those members of $\mathbb{N}_\infty$ or equal to 1 as used in (11.1.5), then the standard part operator can now be applied to (11.1.6) and the result is the same as $\mathbf{st}(\sum_{i=1}^{\lambda} (1/(10^\omega)))$.

The process outlined in (11.1.6) is then applied to finitely many distinct intermediate subparticles — those that characterize an elementary particle. The result is a subparticle each coordinate of which is infinitely close to the value of a numerical characterization or an infinitesimal. When the standard part operator is applied under the usual coordinatewise procedure, the coordinates are either the specific real coordinatewise characterizations or zero. Therefore, N-world formation of particles, the dense substratum field, or even gross matter may be accomplished by a ultrafinite combination of ultrasubparticles that leads to the intermediate subparticle; followed by finite combinations of intermediate subparticles that produce the N-world objects. Please note, however, that prior to application of the standard part operator such subparticles retain infinitesimal nonzero coordinate characterizations in other noncharacterizing positions. (See note (1) on 59.)

We must always keep in mind the hypothesis law [9] and avoid unwarranted speculation. We do not speculate whether or not the formed particles have point-like or “spread out” properties within our space-time environment. These additional concepts may be pure catalyst type statements within some standard N-world theory and could have no significance for either the N-world or NSP-world.

With respect to field effects, the cardinality of the set of all ultrasubparticles clearly implies that there can be ultrafinite combinations of ultrasubparticles “located” at every “point” of any finite dimensional continuum. Thus the field effects yielded by subparticles may present a completely dense continuum type of pattern within the N-world environment although from the monadic viewpoint this is not necessarily how they “appear” within the NSP-world.

There are many scenarios for quantum transitions if such occur in objective reality. The simplest is a re-ultrafinite combination of the ultrasubparticles present within the different objects. However, it is also possible that this is not the case and, depending upon the preparation or scenario, the so-called “conservation” laws do not hold in the N-world.

As an example, the neutrino could be a complete fiction, only endorsed as a type of catalyst to force certain laws to hold under a particular scenario. Consider the set of sentences

$$\begin{aligned} G_B = \{ & \text{An} || \text{elementary} || \text{particle} || k'(i', j') || \text{with} || \\ & \text{total} || \text{energy} || c' + n'. \mid ((i, j, n) \in \\ & \mathbb{N}^+ \times \mathbb{N}^+ \times \mathbb{N}^+) \wedge (1 \leq k \leq m) \}. \end{aligned} \quad (11.1.7)$$

It is claimed by many individuals that such objects as being described in G_B exist in objective reality. Indeed, certain well-known scenarios for a possible cosmology require, at least, one “particle” to be characterized by such a collection G_B . By the usual method, these statements are *- transferred to

$$\begin{aligned} G'_B = \{ & \text{An} || \text{elementary} || \text{particle} || k'(i', j') || \text{with} || \\ & \text{total} || \text{energy} || c' + n'. \mid ((i, j, n) \in \\ & {}^*\mathbb{N}^+ \times {}^*\mathbb{N}^+ \times {}^*\mathbb{N}^+) \wedge (1 \leq k \leq m) \}. \end{aligned} \quad (11.1.8)$$

Hence, letting $n \in \mathbb{N}_\infty$ then various “infinite” NSP-world energies emerge from our procedures. With respect to the total energy coordinate(s), ultrasubparticles may also be ultrafinitely combined to produce such possibilities. Let $\lambda = 10^{2\omega}$ [resp. $\lambda = \omega^2$] and $\omega \in \mathbb{N}_\infty$. Then

$$\sum_{n=1}^{\lambda} \frac{1}{10^\omega} = 10^\omega \text{ [resp. } \sum_{n=1}^{\lambda} \frac{1}{\omega} = \omega \text{]} \in \mathbb{N}_\infty. \quad (11.1.9)$$

Of course, these numerical characterizations are external to the N-world. Various distinct “infinite” qualities can exist rationally in the NSP-world without altering our interpretation techniques. The behavior of the infinite hypernatural numbers is very interesting when considered as a model for NSP-world behavior. A transfer of finite energy, momentum and, indeed, all other N-world characterizing quantities, back and forth, between these two worlds is clearly possible without destroying NSP-world infinite conservation concepts.

Further, observe that various intermediate subparticles carrying nearstandard coordinate values could be present at nearstandard space-time coordinates, and application of the continuous and external standard part operator would produce an apparent not conserved N-world effect. These concepts will be considered anew when we discuss the Bell inequality.

Previously, ultrawords were obtained by application of certain concurrent relations. Actually, basic ultrawords exist in any elementary nonstandard superstructure model, as will now be established for the general paradigm.

Referring back to G_A equation (11.1.1), for some fixed k , $1 \leq k \leq m$, let $h_k: \mathbb{N}^+ \times \mathbb{N}^+ \times \mathbb{N}^+ \rightarrow G_A$ be defined as follows: $h_k(i, j, n) = \text{An}||\text{elementary}||\text{particle}||k'(i', j')||\text{with}||\text{total}||\text{energy}||c'+1/(n')$. Since the set $F(\mathbb{N}^+ \times \mathbb{N}^+ \times \mathbb{N}^+)$ is denumerable, there exists a bijection $H: \mathbb{N} \rightarrow F(\mathbb{N}^+ \times \mathbb{N}^+ \times \mathbb{N}^+)$. For each $1 \leq \lambda \in \mathbb{N}$ and fixed $i, n \in \mathbb{N}^+$, let $G_A(\lambda) = \{\text{An}||\text{elementary}||\text{particle}||k'(i', j')||\text{with}||\text{total}||\text{energy}||c'+1/(n') \mid (1 \leq j \leq \lambda) \wedge (j \in \mathbb{N}^+)\}$. Let $p \in \mathbb{N}$. If $|H(p)| \geq 2$, define finite $M(h_k[H(p)]) = \{A_1||\text{and}||A_2||\text{and}||\dots||\text{and}||A_m\}$, where $A_j \in h_k[H(p)]$, $m = |H(p)|$. If $|H(p)| \leq 1$, then define $M(h_k[H(p)]) = \emptyset$. Let $M^0 = \bigcup \{M(h_k[H(p)]) \mid p \in \mathbb{N}\}$. Please note that the k' represents that “type” or name of the elementary particle, assuming that only finitely many different types exist, i' is reserved for other purposes, and the j' the number of such elementary particles of type k' .

Theorem 11.1.3 *For any $i, n, \lambda \in {}^*\mathbb{N}^+$, such that $2 \leq \lambda$, there exists $w \in {}^*\mathbf{M}^0 - G_A$, ${}^*G_A(\lambda) \subset {}^*\mathbf{S}(\{w\})$ and if $A \in {}^*G_A - {}^*G_A(\lambda)$, then $A \notin {}^*\mathbf{S}(\{w\})$.*

Proof. Let $i, j, \lambda \in \mathbb{N}^+$ and $2 \leq \lambda$. Then there exists some $r \in \mathbb{N}$ such that $h_k[H(r)] = G_A(\lambda)$. From the construction of M^0 , there exists some $r' \in \mathbb{N}$ such that $w(r') = \text{An}||\text{elementary}||\text{particle}||k'(i', 1')||\text{with}||\text{total}||\text{energy}||c'+1/(n')||\text{and}||\text{An}||\text{elementary}||\text{particle}||k'(i', 2')||\text{with}||\text{total}||\text{energy}||c'+1/(n')||\text{and}||\dots||\text{and}||\text{An}||\text{elementary}||\text{particle}||k'(i', \lambda')||\text{with}||\text{total}||\text{energy}||c'+1/(n') \in M[h_k[H(r)]]$. Note that $w(r') \notin G_A$, $h_k[H(r')] \subset \mathbf{S}(\{w(r')\})$ and if $A \in G_A - h_k[M(r)]$, then $A \notin \mathbf{S}(\{w(r')\})$. The result follows by our embedding and $*$ -transfer. ■

The ultrawords utilized to generate various subparticles, whether obtained as in Theorem 11.1.3 or by concurrent relations, are called *ultramixtures* due to their applications. The ultrafinite choice operator $\mathbf{C}_1$ can select them, prior to application of ${}^*\mathbf{S}$. Moreover, application of the ultrafinite combination operator entails a specific intermediate subparticle with the appropriate nearstandard coordinate characterizations. Please notice that the same type of sentence collections may be employed to infinitesimalize all other quantities, although the sentences need not have meaning for certain popular N-world theories. Simply because substitution of the word “charge” for “energy” in the above sentences G_A does not yield a particular modern theory description, it does yield the infinitesimal charge concept prevalent in many older classical theories.

Using such altered G_A statements, one shows that there does exist ultramixtures w_i for each intermediate subparticle and, thus, a single ultimate ultramixture w such that ${}^*\mathbf{S}(\{w_i\}) \subset {}^*\mathbf{S}(\{w\})$. Each particle or elementary particle may, thus, be assumed to originate from w through application of the ultralogic ${}^*\mathbf{S}$.

11.1.2 More on Subparticles

The general process for construction of all N-world fundamental entities from subparticles can be improved upon or achieved in an alternate fashion. One of the basic assumptions of subatomic physics is that in the Natural-world two fundamental subatomic objects, such as two electrons, cannot be differentiated one from another by any of its Natural-world properties. One of the conclusions of what comes next is that in the NSP-world this need not be the case. Of course, this can also be considered but an auxiliary result and need have no applications. At a particular instant of (universal) time, it is possible to associate with each entity a distinct “name” or identifier through

subparticle construction. This is done through application of the subparticle naming coordinate a_1 . As will be shown, the concept of independent *-finite coordinate summation followed by n-tuple vector addition can be accomplished by means of a simple linear transformation. However, by doing so, the concept of the *-finite combinations or the gathering together of subparticles as a NSP-world physical-like process is suppressed. Further, a simple method to identify each N-world entity or Natural-system is not apparent. Thus, we first keep the above two processes so as to adjoin to each entity constructed an appropriate identifier.

The basic construction may need only an infinite *-finite “number” of subparticles to construct ALL of the fundamental subatomic or field constituents, if they exist, for a universe. If we were to allow denumerably many fundamental entities, as will be seen, we would still have, at the least, a *-denumerable set of subparticles from which to construct “other things,” whatever they may be. From the general paradigm approach, the set of all ultrasubparticles, at the least, has the same cardinality as the hypernatural numbers, $^*\mathbb{N}$. We now begin a more specific construction. First, we assume that there is but a finite set of physical properties that characterize the fundamental entities and that distinguish the fundamental entities one from another. These are the accepted entities from which the constituents of a universe are assumed to be constructed. Let natural number $f \geq 1$ denote this finite number. [This need not be the fundamental entities that, as of today, are deemed as correct by the particle physicists.] What is also important in this construction is that according to the discussion in the paragraph prior to theorem 11.1.1,

(1) for each positive non-negative real number r there exists a unique $\lambda_r \in \mathbb{N}_\infty$ such that $\text{st}(\lambda_r/10^\omega) = r$. For $r = 0$, use $\lambda_0 = 0$. This gives an injection $N: \mathbb{R}^{\geq 0} \rightarrow (\mathbb{N}_\infty \cup \{0\})$. [Note that in practice the 0 is not needed since the standard part of any of the original infinitesimal coordinate values is zero.] Further, the fact that the inverse N^{-1} exists is of great importance. Notice that the nonzero λ_r s are members of distinct mathematical galaxies. Further, if the standard part operator is restricted to this unique family of λ_r s, then the map $\text{st}^{-1}(r) = \lambda_r/10^\omega$ is an appropriate inverse.

(2) For the set of all fundamental entities, there is a maximum finite number f of $\lambda_r \in \mathbb{N}_\infty$ used, under the concepts of independent coordinate summation and the standard part operator, to characterize fully each entity.

Let $P \subset \mathbb{N}$ be the infinite set of all prime numbers. In general, there is a prime number H such that the cardinality of $\{x \mid (x \in P) \wedge (2 \leq x < H)\} \geq f$. This can be re-expressed in terms of maps. Thus, there is an injection $j: [1, f] \rightarrow \{x \mid (x \in P) \wedge (2 \leq x < H)\}$. We know that this injection can be replaced by a similarity map (i.e. order preserving bijection) J from $[1, f]$ onto the range of j which is a subset of $\{x \mid (x \in P) \wedge (2 \leq x < H)\}$. Hence by *-transfer, the internal similarity map $^*J: [1, f] \rightarrow \{x \mid (x \in ^*P) \wedge (2 \leq x < H)\}$. Since $f \geq 1$ is a standard number, then the set $^*J[[1, f]]$ is a finite internal and standard set. What we now intend to do is a “mini” form of Gödel style encoding. First, consider the internal naming set $^*K = ^*P - ^*J[[1, f]]$. This set of *-primes is *-denumerable. [For simplicity and for comprehension, one can consider the set $^*J[[1, f]]$ as the first f standard primes numbers in order.] The internal set *K is corresponded to a (internal) set of ultrasubparticles that can be used to construct ALL of the fundamental entities used to construct ALL of the constituents of a Natural universe. This is done by considering the ultrasubparticles to be modeled by *-finite n-tuples as they are represented by of *-finite sequences. In what follows, we’ll be somewhat less formal in our construction.

(3) Let $^*\mathbb{R}^{[1, n]}$, where $n \in \mathbb{N}_\infty$, denote the internal set of all internal functions defined on $[1, n]$ into $^*\mathbb{R}$. We now consider the set of ultrasubparticles as modeled by all *-finite sequences with the

following values. For $i = 1, a_i = k$, where $k \in {}^*K$, the naming coordinate. For $i = 2, a_i = 1$, the counting coordinate. For $i \geq 3$ and $*$ -odd, let $a_i = 10^{-\omega}$. For $i \geq 4$ and $*$ -even, let $a_i = -10^\omega$. [Remember that fixed $\omega \in \mathbb{N}_\infty$.] The internal set $\mathcal{U}$ of all such $*$ -finite sequences (n-tuples) models the collection of all ultrasubparticles needed to construct ALL of the constituents that comprise a Natural universe.

Note that every member of $\mathcal{U}$ has a distinct identifiable name in the naming coordinate.

(4) Let $\lambda_r \in {}^*\mathbb{N}_\infty$ be used to obtain one intermediate subparticle. First, the specific identifier of this intermediate subparticle is obtained by *multiplying* the $*$ -prime identifiers of the individual ultrasubparticles.

(5) Then let $J(i)$ be the identifier for the specific characteristic associated with this specific λ_r . Now consider $J(i)^{\lambda_r}$ as multiplied to the $*$ -finite natural number obtained in (4). This is the complete identifier used in the a_1 place for each intermediate subparticle. Further, the single $*$ -finite n-tuple used to model this intermediate subparticle has $a_2 = \lambda_r$ and $a_{J(i)} = \lambda_r/10^\omega$ or $a_{J(i)} = -\lambda_r/10^\omega$ and all coordinates defined as above or as in (3).

(6) A fundamental entity is obtained by combining together a finite set of intermediate subparticles. This is modeled by the usual n-tuple addition with respect to the finitely many intermediate subparticles that comprise a fundamental entity with one exception. The a_1 is obtained not by coordinate addition but by multiplication of the finitely many intermediate a_1 coordinates.

(7) By the fundamental theorem of number theory $*$ -transferred to ${}^*\mathbb{N}$, the coordinate a_1 obtained in (6) is unique in the following sense.

(a) Each and every fundamental entity used to construct any composite object has a different identifier.

(b) For each fundamental entity, in the a_1 coordinate representation, the finite set of factors $J(i)^{\lambda_r}, \dots, J(j)^{\lambda_s}$, where $i, j \in [1, f]$, identifies the type of fundamental entity. The remaining prime factors identify each and every one of the distinct ultrasubparticles the are used to obtain each and every fundamental entity.

(c) Although the $J(i)^{\lambda_r}, \dots, J(j)^{\lambda_s}$ may be identical for two or more fundamental entities, the natural number identifier composed of the remaining factors would yield distinct identifiers for each of the required Natural-world physically identical fundamental entities. These distinct identifiers could only be accounted for within the pure NSP-world.

(8) When an intermediate subparticle is obtained by independent coordinate summation, this same summation is applied to the counting coordinate a_2 . Hence, for a specific intermediate subparticle obtained for a specific λ_r , the $a_2 = \lambda_r$. Consequently, the n-tuple that models a specific fundamental entity has as its a_2 coordinate the finite sum of the necessary λ s. This number is a member of $\mathbb{N}_\infty$ and represents the number of specific ultrasubparticles used to construct a specific fundamental entity.

(9) Observe that the 0 is never used to describe a specific fundamental entity characteristic. Thus for things such as a standard photon, we do not state the characterization that it has no rest mass. The fact that the coordinate that represents rest mass is infinitesimal is interpreted to mean that it does not have a rest mass. The actual Natural-world behavior of a fundamental entity is modeled by the n-tuple obtained by applying the standard part operator only to the finitely many a_i that are non-infinitesimal limited (i.e. finite, bounded) $*$ -reals.

(10) Notice that our notation for independent coordinate summation can be expressed as follows:

$$\sum_{n=1}^{\lambda} 1/10^{\omega} = \sum_{n=1}^{\lambda} 1^n/10^{\omega} \text{ or}$$

$$\sum_{n=1}^{\lambda} -1/10^{\omega} = \sum_{n=1}^{\lambda} -(1^n)/10^{\omega}.$$

Due to the uniqueness of all the identifiers and the procedures followed to obtain a specific fundamental entity, there are obvious inverse procedures that can be applied to obtain all of the original ultrasubparticles used to construct a specific fundamental entity. For example, all of the construction features can be obtained by examining the identifier a_1 for a specific fundamental entity.

What is needed is that we show that there are “enough” objects in $\mathcal{U}$ to construct a Natural universe and that we will have a “great many” left over, if we wish, for other purposes. We note that $K = P - J[[1, f]]$ is denumerable. Consider any partition $\mathcal{Q}$ of K into nonempty finite sets. The infinite set $\mathcal{Q}$ is a subset of the set of all finite subsets $\mathcal{K}$ of K . But $\mathcal{K}$ is denumerable. Hence $\mathcal{Q}$ is denumerable. But the naming portion of a_1 for each fundamental entity is but a distinct member of ${}^*\mathcal{K}$. Hence, the (external) cardinality of ${}^*\mathcal{K}$ determines the “number of” fundamental entities that can be used to construct the composite objects for any Natural universe. It is well known that the external cardinality of ${}^*\mathcal{K}$ is greater than or equal to the degree of saturation of the nonstandard model being considered. Thus the cardinality of the set of fundamental entities can be controlled to such an extent that even if one were to assume that a Natural universe contains at the most a specific infinite cardinal number of composite objects constructed from finitely or even a specific maximum infinite cardinal number of fundamental entities, then it is possible to consider a nonstandard model saturated to such a degree that there are enough fundamental entities to construct such a Natural universe. Further, the cardinality can be increased to any level that will allow for subparticles to be used within the NSP-world itself for such things as ultrafast informational transmissions “instantaneously ” throughout the entire Natural universe.

Notice that the unique naming coordinate for each fundamental entity also yields a unique name for each Natural object or Natural-system that is formed or that is altered in anyway. This allows for a form of complete control over all such objects as well as allowing instantaneous informational subparticles to be specifically directed to specific entities or combination of entities.

The process of applying the standard part operator does not erase the original coordinates of the constructed objects. The process only yields the N-world measurements. Now alteration of the fundamental subatomic or field entities is modeled by vector space subtraction and this would be followed by independent *-finite subtraction. This represents a “breaking apart” of the fundamental entity into its original ultrasubparticle constituents. The original identifiers are obtained since they are but the non-“f” prime factors that appear in the identifier of the constructed fundamental entity or Natural-system. An altered constituent is then obtained by repeating the construction process. This can be used to eliminate the virtual particle or process concept within reality models. The models that need such concepts to predict behavior are now just that “models” for predicted behavior and would not require the virtual “stuff” to exist in reality.

As mentioned, the formation of fundamental entities from ultrasubparticles through the process of the intermediate subparticle formation can be modeled by means of a very simply (diagonal styled) *-continuous linear transformation. Let a fundamental entity have the required finitely

many coordinate numerical characteristics as discussed previously and let nonempty finite $A \subset {}^*\mathbb{N}$ be the specific coordinate names. Then there is a finite set of hypernatural numbers $\{\lambda_i \mid i \in A\}$ that represent the number of summands in the respective $*$ -finite summation process. (The process that yields the intermediate subparticles.) Consider an $n \times n$ hypermatrix with the diagonal elements $\lambda_i = b_{ii}$, $i \in A$, and $b_{jj} = 1$, $j \in ({}^*\mathbb{N} - A)$, $j \leq n$ and $b_{ij} = 0$ otherwise. Then letting (a_j) be a coordinate representation for an ultrasubparticle, consider $(b_{ij})(a_j)^T$. This hypermatrix represents an internal function that will take a single *representation* for an ultrasubparticle and will yield, after application of the standard part operator to the appropriate coordinate values (or all of them if one wishes), a *representation* for a fundamental entity. BUT note that identifier coordinate would not be that of the intermediate subparticle. Unless we use the sum of the λ_i s in the second diagonal place, the counting coordinate would not be the appropriate value. Of course, the matrix inverse would yield a representation of an original ultrasubparticle with its identifier. Thus using either method, the identifier would be lost to the N-world. But using one of the favorite simple mathematical models, the matrix approach, the fundamental entity identifier would be lost to the NSP-world as well. This shows how two different mathematical procedures can lead to equivalent N-world results, but yield considerably different NSP-world ramifications.

The next step in a subparticle re-formulation of particle physics would be to consider specific NSP-world predictions for what is claimed are random or, at least, probabilistic in character actual observed events. In this regard, I mention the known fact that a probability density functions behavior can be predicted by a specific well-defined sequence. For example, it is known that there is a natural number x and a specifically defined sequence $x2^n$ such that if one takes the fractional part of each $x2^n$, less than $1/2$ and correspond it to H, and the fractional part greater or equal to $1/2$ and correspond it to a T, then the predictive results will pass every statistical test that implies that H and T are randomly obtained by a sequence of H = heads, T = tails coin tosses. Clearly, this process, as far as the sequence $x2^n$ is concerned is a predictable process and contradicts scientific probabilism. But the specific numbers x need not be known by entities within the Natural universe.

11.2 An Ultimate GGU-model Conclusion.

In the book “Ultralogics and More” other subparticle properties are discussed. These include the ultraenergetic and ultrafast subparticles. The ultrafast subparticles are of significance in that they can be used to provide seemingly instantaneous informational transmissions throughout the Natural universe. This is significant for the ultralogic and ultraword generation of a universe. Clearly the research presented in this monograph is not complete but is only designed to present the beginning concepts in what is hoped will be a continuing research activity. One important question is relative to the fact that an ultimate ultraword such as w' generates a preselected or theory generated ideal universe. How can this be made to correspond to the actual universe in which we live where each Natural-system is perturbed or altered within certain limits dictated by Natural law from this ideal case? Such alterations can also be produced by partially independent agencies such as biological entities. There are different ways to attack this problem but the existence of the UN-events would require that speculation be restrained.

One method is slightly similar to the Everett-Wheeler-Graham many-worlds interpretation (parallel universes) but is much less esoteric in character. One can consider countably many w'_t from Corollary 7.3.4.1 each one containing the allowable alterations in Natural-system behavior starting from the moment t . Denote the corresponding set of universes by $\mathcal{U}$. At any instant of substratum time t , all the universes but one are “covirtual” universes. This does not necessary mean that they

actually exist in some type of NSP-world reality. This simply means that they exist potentially. When any one or more Natural-systems is perturbed then the UN-events react first so to speak. Information relative to this reaction is transmitted within $\{w'_t\}$ by means of ultrafast subparticles. This information causes a w'_{U_t} to be selected such that $U_t \subset {}^*\mathbf{S}(\{w'_{U_t}\})$, $U_t \in \mathcal{U}$. For the “next” instant this generated U_t becomes the Natural universe objective reality. Now technically there would be a “next” instant since there would be an NSP-world time interval over which all of the Natural-systems would have a fixed frozen segment segments that would not be altered. The only possible alterations would be in the UN-events. This process then continues throughout all of Natural universe time.

There is a second approach that can solve this perturbation problem but requires a complete restructuring of ultraword generation. This method is similar to the concept of “parallel” logic. Time in this case is modeled as previously done but on interval $(-\infty, +\infty)$. At a particular observational beginning instant t , a time slice in the mathematical sense is obtained. This slice will yield a frozen segment describing a Natural event for each of the present Natural-systems. This will lead as in Chapter 7 to an ultraword x that will by means of ${}^*\mathbf{S}$ generate this slice. Let $\mathcal{NA}$ denote the set of all potential Natural-systems. This means that some of these at the instant t contain only frozen segments from ${}^*\mathbf{T}_i - \mathbf{T}_i$.

Assume that $\mathcal{NA}$ is at the least denumerable. Then each x_t (the ultraword associated with the instant t) also contains UN-events. Again the alterations in Natural-system behavior within the limits of Natural law lead to changes in the frozen segment descriptions for the very “next” Natural event. These changes are incorporated within the NSP-world time interval into the “next” ultraword x_{t_1} . Application of ${}^*\mathbf{S}$ to x_{t_1} now yields the “next” slice. This process continues throughout all of the Natural universe time frame. Of course the “time” being referred to here is not N-world measured time but the time concept previously discussed that is concerned with “before and after.”

The first of the above models appears to be the most significant and the one that is probably the closest to objective reality. This portion of this model leads to the moment-to-moment re-generation of our universe since its behavior most certainly is perturbed moment-to-moment from the ideal.

CHAPTER 11 REFERENCES

- 1 Beltrametti E. G. Enrico G. and G. Cassinelli *The Logic of Quantum Mechanics* Encyclopedia of Mathematics and Its Applications Vol. 15 Addison-Wesley Reading 1981.
- 2 d’Espagnat B. The quantum theory and realism Scientific America 241(5)(1979) 177.
- 3 *Ibid.*
- 4 *Ibid.* 181.
- 5 *Ibid.*
- 6 *Ibid.* 180.
- 7 Feinberg G. Possibility of faster-than-light particles Physical Review 159(5)(1976) 1089–1105.
- 8 Hanson W. C. The isomorphism property in nonstandard analysis and its use in the theory of Banach Spaces J. of Symbolic Logic 39(4)(1974) 717–731.
- 9 Herrmann R. A. D-world evidence C.R.S. Quarterly 23(1986) 47–53.
- 10 Herrmann R. A. The Q-topology Whyburn type filters and the cluster set map Proceedings Amer. Math. Soc. 59(1)(1975) 161–166.
- 11 Kleene S. C. *Introduction to Metamathematics* D. Van Nostrand Co. Princeton 1950.

- 12** Prokhovnik S. J. *The Logic of Special Relativity* Cambridge University Press Cambridge 1967.
- 13** Stroyan K. D. and W. A. J. Luxemburg *Introduction to the Theory of Infinitesimals* Academic Press New York 1976.
- 14** Tarski A. *Logic Semantics Metamathematics* Clarendon Press Oxford 1969.
- 15** Thurber J. K. and J. Katz Applications of fractional powers of delta functions *Victoria Symposium on Nonstandard Analysis* Springer-Verlag New York 1974.
- 16** Zakon E. Remarks on the nonstandard real axis *Applications of Model Theory to Algebra Analysis and Probability* Holt Rinehart and Winston New York 1969.
- 17** Note that the NSP-world model is not a local hidden variable theory.
- 18** Herrmann R. A. Fractals and ultrasmooth microeffects J. Math. Physics 30(4) April 1989 805–808.
- 19** Davis M. *Applied Nonstandard Analysis* John Wiley & Sons New York 1977.

(1) For subparticles, only two possible intrinsic properties for elementary particle formation are here considered. Assuming that there are such things as particles or elementary particles, then they would be differentiated one from the other by their intrinsic properties that are encoded within subparticle coordinates. When there are particle interactions, these intrinsic properties can be altered or even changed to extrinsic properties. How the alteration from intrinsic to extrinsic occurs probable cannot be known since it most likely is an ultranatural event. For new results on subparticle formations, see <http://arxiv.org/abs/quant-ph/9909078>

GLOSSARY

ANAMORPHOSIS PROCESS. A process that takes place after the intermediate or minimum metamorphic effect. This process distorts the original electromagnetic or particle radiation so that observations taken after the time fracture cannot give any information about the condition of our universe prior to the time fracture.

DEDUCTIVE WORLD MODEL (D-WORLD MODEL). This is the mathematical structure, the extended G-structure, interpreted linguistically.

DEVELOPING NATURAL SYSTEM. A natural system that is altered in some identifiable manner with respect to time.

DEVELOPMENTAL PARADIGM. This is a sequence of broadly defined descriptions for the moment-to-moment behavior of a Natural-system. It corresponds to an event sequence.

EFFECT. A described and named phenomenon, usually Natural or ultranatural, that is closely associated with a specific cause. It is the claimed result of the cause. As with the concept of a cause, what is considered as an effect of a cause need not be based upon any verification, especially where other effects not so accepted can be considered as resulting from the same cause. The symbol string that names an effect is taken from a list symbol strings categorized as effects.

EVENT. For this book, this is a Natural or ultranatural phenomenon that is being described by a string of symbols.

EVENT SEQUENCE. A time ordered collection of events.

GGU-MODEL. An abbreviation for the phrase “General Grand Unification Model.” This is a collection of statements written in a physical-like language that gives a description for processes and

objects that exist logically in the NSP-world and that produce universes or alter their development. The GGU-model can also be used for other purposes.

GRUNLEGEND STRUCTURE (G-STRUCTURE). The basic mathematical structure used for both the D-world and GGU-models.

IN-PROCESS. See intrinsic natural process.

INDIRECT VERIFICATION. A portion of the so-called scientific method that claims that the existence of entities and processes that cannot be sensed by human beings directly can be inferred from predicted events that can be directly sensed by human beings or their machines.

INFANT. See subparticle.

INFINITESIMAL. A mathematical entity that, in this paper, is used to measure physical behavior. Originally, such measures were thought to exist in reality. Then, when contradictions occurred, although still used, they became physical parameters or catalysts. But, today, since the contradictions have been removed, they can once again be interpreted as measures for real physical entities.

INFORMATIONAL TRANSMISSIONS. These occur when special Natural or ultranatural entities are used to give certain special information to other Natural or ultranatural entities. The methods by which such information is shared may be unknown.

INTRINSIC NATURAL PROCESS. Natural processes that are hidden from direct verification and that mirror certain specific human mental processes.

INTRINSIC ULTRANATURAL PROCESS. These are processes within the pure NSP-world that when restricted to the Natural world become intrinsic Natural processes.

INTRINSIC ULTRANATURAL SELECTION. This is the process within the NSP-world that when restricted to the Natural world mirrors the human procedure of selecting finitely many objects from a collection of objects.

IUN-PROCESS. An abbreviation for the phrase “intrinsic ultranatural process.”

IUN-SELECTION. An abbreviation for the phrase “intrinsic ultranatural selection.”

LINGUISTIC INTERPRETATION. A correspondence between certain processes and objects used within or associated with the science of linguistics, and a special mathematical structure - the extended G-structure.

LOGIC COMPUTER. A computer that can take a finite collection of statements and using a specific program (i.e. set of rules) produce the same written statements that can be produced by the human mind using the same set of rules.

METAMORPHIC. This means a special type of sudden or instantaneous change where such a change is instantaneous only from a Natural world viewpoint.

N-EVENT. An abbreviation for the phrase “Natural event.”

N-OBJECT. An abbreviation for the phrase “Natural object.”

N-PROCESS. An abbreviation for the phrase “Natural process.”

NATURAL EVENT. An event that is assumed to take place within our universe.

NATURAL INITIAL CONDITIONS. Described conditions that must occur (or must be met) before a Natural event can occur. Initial conditions also can differentiate one Natural event from another distinct Natural event.

NONSTANDARD PHYSICAL WORLD MODEL. This is a theory that uses physical-like terms in such a way that each of the physical-like terms corresponds to a term within a mathematical theory. The term nonstandard refers to the special mathematical theory to which the physical-like terms correspond. The GGU-model is a portion of this theory.

NSP-WORLD MODEL. An abbreviation for the phrase “nonstandard physical world model.”

OBJECTIVE REALITY. This is the assumed reality that exists outside of the mind and, thus, outside of the imagination.

PARAMETER. Can take on various meanings. It can be a catalyst or a quantity which must be known and inserted into a physical theory before that theory will predict Natural-system behavior specifically.

PHYSICAL INTERPRETATION. Taking data gathered by human beings or machines and associating this data with terms taken from a language that employs terms called physical terms. Physical terms are often assumed to name processes and entities within our universe. This also means the construction of a mathematical model by associating physical terms with abstract mathematical entities.

PHYSICAL THEORY. All of the logically deducible results that one can obtain from a fixed set of statements written in physical terms.

PREDICTION. Another name for a deductive conclusion obtained from physical hypotheses. Sometime, a conclusion within a physical theory that might be directly measured within a laboratory setting.

PURE NSP-WORLD. This is composed of all things that are in the NSP-world and not in the Natural world.

REALISM. The selection of entities within a mathematical theory as representing objects that are claimed to exist in reality. Absolute realism is the philosophy that all entities within a mathematical structure correspond to real physical objects.

REDUCTIONISM. The philosophy that all, microscopic, macroscopic and large scale physical behavior is produced from a finite collection of physical objects and fundamental interactions that exist and operate on the subatomic microscopic level.

STANDARD MODEL. This is the Big Band model although there are some variations.

STRING OF SYMBOLS. Basically these are but marks that are written down on any appropriate surface. Like the marks that appear between the B and the . of the previous sentence. However, today, this would include almost all forms of human perception since such marks can lead to the replication of much of what we perceive with our senses.

SUBJECTIVE. Within the mind or imagination.

SUBNATURAL-SYSTEM. The same meaning as a Natural-system except that this type of Natural-system is recognized as a portion of another Natural system.

SUBPARTICLE. A pure NSP-world object that can be shown to exist logically. They have many properties that can be described and many properties that cannot be described. They should not be thought of as particles, however. A better name would be “things.”

SUBSTRATUM UNIVERSE. The same as a background universe. In general, the term substratum means a declared basic foundation that is contained in a “larger” something. In this case,

this is a basic foundation for universes and, contrary to the general meaning of substratum, the universes are contained in the substratum universe.

SUDDEN CHANGE. Also called abrupt change. These are changes that occur in the behavior in a Natural-system that appear to be discontinuous in character. They may appear to occur instantaneously. The change need not be sudden as viewed from the NSP-world.

TECHNICAL LANGUAGE. A language that one learns when one studies a distinct discipline and used for the special concepts within that discipline.

THEORY. This is the entire set of conclusions that can be deduced logically from a set of premises.

TIME FRACTURE. A specific moment during the development of a Natural system when Natural time, if it existed prior to this time, is suspended and processes within the NSP-world alter the development of such a Natural system. (Note that such a specific moment can also correspond to a specific temperature or other similar parameter.) This NSP-world process takes place during allowable NSP-world time. If there is no Natural time reference for the fracture, then this fracture is called an ultra-fracture and if it is within an ordered sequence of events, then it corresponds to a specific universal event number.

THING. See subparticle.

ULTIMATE ULTRAMIXTURE. This is the ultraword that generates by application of an ultralogic a particle or elementary particle.

ULTIMATE ULTRAWORD. A single object that exists logically within the pure NSP-world and that is used to form all of the Natural-systems that comprise an entire universe.

ULTRA. A prefix used to distinguish NSP-world processes and objects from similar sounding Natural world processes or objects. However, each of the objects or processes so distinguished does have one or more properties in common with the process or object named by the term following the prefix ultra.

ULTRAFAST. This is a characterization associated with certain pure NSP-world subparticles. This characterization indicates that such subparticles can traverse, within the NSP-world, from one point within our universe to any other point within our universe without expending any Natural time.

ULTRALOGIC. This is the force-like pure NSP-world process that when applied to an ultraword produces any Natural-system including an entire universe. It also produces subparticles.

ULTRAMIXTURE. This is the ultraword that generates by application an ultralogic an intermediate subparticle.

ULTRANATURAL EVENT. An event that takes place within the pure NSP-world. This event, usually, effects indirectly Natural events. Detailed information about such events cannot be described in any language used by any entity within the Natural universe.

ULTRANATURAL INITIAL CONDITIONS. The same as Natural initial conditions, but these conditions must be met within the pure NSP-world.

ULTRANATURAL LAWS. This term has the same meaning as the expression Natural laws. But these fixed processes take place within the pure NSP-world.

ULTRANATURAL OBJECT. An object which is contained in the pure NSP-world.

ULTRASUBPARTICLE. These are entities within the GGU-model that are essentially of a single type. These entities can be used to construct all of the fundamental constituents of our universe, including “empty” space, by means of a simple finite-like “gathering together” process.

ULTRASMOOTH. A process that can be characterized by the smooth pattern of the curved portion of the diagram . However, this process would still appear to have such a smooth pattern even if view through a microscope of infinite power.

ULTRAUNIFORM. An extremely fine step-by-step process. So fine that it would not appear to be broken into disjoint pieces even if viewed through a microscope of infinite power.

ULTRAWORD. An unusual object that exists logically within the pure NSP-world and contains all of the necessary concepts, laws, building plans and the like that control the formation and the properties of a specific Natural-system. It is this object that a specific ultralogic acts upon to produce a Natural-system.

UN-EVENT. Abbreviation for the phrase “ultranatural event.”

UN-LAWS. An abbreviation for the phrase “ultranatural laws.”

UN-OBJECT. An abbreviation for the phrase “ultranatural object.”

VIRTUAL (PARTICLE OR PROCESS). These are subatomic objects or processes that many quantum physicists believe exist in reality that mediate interactions between other objects but are not themselves detectable due to the time span over which they exist. Or they can simply be members of a list that states that such and such object or process is virtual relative to a specific interaction. (There are other more technical definitions.)

grand 2, 3, 4, 5.

Corections made 27 AUG 2005, Section 11.1.2 added 12 APR 1995, Section 11.2 added 24 DEC 1996

E-mail post of 12 OCT 1994.

Since sending out, to members of this network, requested copies of my GGU-model solutions paper, certain questions have been asked of me. I will provide answers to the more significant questions to all network individuals that have by my records been supplied with this paper.

(1) Is there a strong argument that there can be NO bottom-up standard unification?

Yes, there is. Under the usual hypotheses associated with the various quantum theories, it is claimed that foundation for these theories is not classical logic but, rather, is quantum logic. This false claim would produce a direct contradiction since the theory we call quantum logic is produced by means of classical logic. What is meant, but not properly stated, is that the behavior within the microphysical world of QM is not classical Newtonian behavior. On the other hand, the underlying logical basis for macroscopic mathematically based scientific theories is the classical logic AND the behavior of many such natural systems is approximated by classical Newtonian concepts. It would appear that microphysical behavior and classical behavior are incompatible. Obviously, this implies that unless the basic hypotheses are alter for theories that claim to predict natural system behavior, then it appears not to be possible to construct a consistent standard unification for all standard

physical theories. On page 45 and elsewhere of the above mentioned paper, I discuss how the GGU-model approach eliminates this difficulty IF the behavior is viewed from the nonstandard physical world.

Another important aspect of the quantum theory difficulties is illustrated by Feynman diagrams and QED. There exist(ed) the discreteness paradox problem among others. In his Nobel Lecture Feynman stated the he did not want to use the “contact” approach relative to his Feynman diagrams. Because the assumption that fields, virtual particles and virtual processes were considered by his colleagues to exist in reality, but his mathematical approach produced ideas that they did not consider to exist in reality, he said “. . . the idea (of action at a distance) seemed so obvious to me and so elegant that I fell deeply in love with it.” But due to various philosophical requirements, his theory was altered to include all the necessary virtual concepts and especially energy transport by means of photons. Of course, when Einstein introduced the concept of the photon he wrote that it was an imaginary entity. Feynman later (after the Nobel prize), stated that the idea of photons actually doing such a thing was a “crazy” idea.

In a 1983 Nature and System paper, “Mathematical Philosophy and Developmental Processes,” the discreteness paradox is explained and at the same time the beginnings of the theory of sub-particles is explored. (If you wish a copy of this paper, let me know.) The discreteness paradox is the problem on the boarder between quantum theory with its apparent discontinuities such as the “sudden” emission of a photon and continuous theories that predict the exact same outcomes. It is shown in that paper, where the mathematics is contained explicitly in the monograph “Ultralogics and More,” that there is no paradox. Indeed, both theories are but restrictions of various hypercontinuous etc. processes when viewed from the NSP-world. If you couple this resolution of this so-called paradox with subparticle hyperfinite and finite composition processes as partially described on pages 47 – 53 of the “solutions” paper, then all virtual particles and processes and even photon energy transport can be removed from the Feynman diagram interactions. One could return to Feynman’s original belief. The entities within the mathematical model that correspond to such eliminated processes are but parameters or modeling techniques or what are sometimes called metaphors or catalysts – they are not to be considered objectively real in character. To show that the selection of mathematical entities as objectively real or parameter in character is but philosophical in character, ask a physicists whether infinitesimal quantities actually exist in some type of objective reality – quantities needed to predict behavior within most models. Then read the quotation from Robinson on page 6.

Then there is the idea of the point-like behavior of the electron (or maybe one dimensional strings moving in time), yet we have the concept of emission and absorption of photons. [Is there an internal property of a point-like element that can produce an emission or absorption?] Has it ever occurred to the physics community that the philosophy of science DOES NOT INCLUDE the requirement that a MODEL and reality correspond. The definition of a model, in general, is that it *mirrors behavior*. This includes my subparticle theory. When quark theory was first proposed, it was call the Quark Model. And part of the quark model QCD is explicitly said to mirror humanly comprehensible color concepts. In all such models, the appropriate phrase “[They](It) behave(s) as if [they] (it) [are] (is)” may be added without altering anything except the unfounded realism concept. With this alteration, all of the predictions for the actual behavior of which is actually observed by human or machine sensors would still hold. From such an alteration, from substitution of subparticle theory for virtual entities, etc., it follows that, within science, prediction does not imply that such model elements exist in objective reality.

Then it seems that many biologists believe that finding a unifying theory for the so-called four fundamental interactions (forces) will not lead to the production of what they call “emerging properties.”

(2) Does a proper unification need to predict new behavior?

The answer to this question is NO. As put forth by the Wheeler group at Princeton and from the very definition of the concept, a unification of all physical theories is not intended nor need it predict new standard physical behavior. The concept itself was advanced for the one purpose of providing a logical unification of physical theories that do not at present have such a logical connection. Indeed, to unify even those theories that use inner-logical procedures that may not be *n*th-order in character such as intuitionistic logic. Please note that string theory falls into this category since it seems impossible to verify any of the conclusions of string theory by predicting new macroscopic behavior that can be verified within the laboratory. The same is true with the Everett-Wheeler-Graham multi-worlds interpretation for QM.

The Princeton group tried to construct such a unification from the statistics of long propositions from the simple propositional (logic) calculus. This indicates that such a unification is not intended to predict new behavior. Under the guiding rules for such a unification, as listed on pages 5 and 6, the GGU-model is such a unification. As such, it unifies all standard physical theories and predicts all standard observations predicted by such theories and also it produces those physical events for which there is, at present, no standard physical theory. It produces a logical connection between all physical behavior. It does this with a compatible logical process that is, in its standard form, a subsystem of all of the logical systems used within all of the present physical theories. Why doesn't this lead to an inconsistent theory as in the standard case?

This connection is by means of ultrawords and ultralogical operators. There are NO actual (standard) physical hypotheses except for the behavior of logical operators and a few linguistic (standard) procedures. The ultrawords and especially the ultimate ultraword encapsulate the entire consequences and hypotheses of the predicting theories as well as all other ideal non-theory produced behavior and other ultranatural stuff. Since this is the opposite of the claims of the reductionists, they will find it very hard to comprehend unless they are open to a change their underlying philosophy.

The GGU-model does predict the existence of sustaining ultranatural events ultranatural laws etc. The model also states on its own and without any human interpretation that while humankind is in its present mode of existence, humankind cannot describe in any standard human language the specifics associated with these ultranatural events. Further, the GGU-model approach specifically highlights the different philosophical bases of the different scientific methods that have been developed over 2500 years. Not only do you need to decide what entities are only technical constructs for processes that are in reality beyond human comprehension, the model predicts rationally what Planck believed that scientists will not be able to explain fully or comprehend the behavior of all natural systems. Indeed, in modern model building, the necessity for accepting one model over another due to an increased predictability has long been discarded. The most important aspects have been simplicity, the underlying philosophical basis, its explanatory power and, most important of all, the development of strong logical arguments, arguments that do not exist in the standard quantum or relativity theories.

Sincerely,
R. A. Herrmann

A Misunderstood Model

1. Having now read many of the previous posts relative to a possible cyclic universe produced through quantum fluctuations, I am now more convinced than ever that I have done a very poor job of explaining the major properties of the GGU-model. [I point out that the philosophy of such cyclic universes and their possible mechanisms are discussed in the 1974 Wheeler paper “Is Physics Created by a Cosmogony?,” where they are rejected for various reasons, and from which I took the scientific requirements for a pregeometry; requirements met by the general GGU-model.]

2. I should have chosen an entirely different name for this mathematical model since the name has led to unfortunate misunderstandings. I am aware of the fact that there are some members of this CRSnet who are not particularly fond of the GGU-model approach. I apologize for not making many of its more significant aspects more accessible, although they are in reality often rather simple in character. I suggest that this post be retained for further study and reflection for this is not a model just for the sudden appearance or sudden alterations in natural system behavior. It is much, much more than that.

3. From the viewpoint of universe creation, the model gives GENERAL methods for the creation of universes. Please note the plural. The model is itself INDEPENDENT from any actual physical theory for how any universe will sequentially develop (i.e. a specific cosmology). The processes used in the GGU-model are INDEPENDENT from any physical theory for the development of any natural system contained within a universe, although it will generate such behavior from more fundamental constituents. This must be the case for any general processes for the creation of different universes. The model is a COSMOGONY not a cosmology. The model automatically states that there are identical but identifiable entities that have ONE fixed set of properties and that can be used to create infinitely many different universes and not just the one in which we dwell. Indeed, these entities can also model and replace various “vacuums” as well as replace virtual processes or particles. The model also produces the various “laws of nature” needed to drive each of the created natural systems. It has other properties, but what this is saying is that certain universes can be produced.

4. With respect to any particular cosmology, the model solves the GENERAL grand unification problem in that it unifies the behavior of all natural systems that comprise ANY of the produced universes.

5. This model IS LOGICALLY FALSIFIABLE. It does not produce all possible universes. Each universe that it produces must develop with respect to a certain logical pattern. The ultranatural processes, ultranatural events, ultra-anything are NOT POSTULATED, but they are PREDICTED by modeling the simplest aspects of human thought, written communication, and selection. Sub-particles are NOT POSTULATED, but they are PREDICTED from one of the basic properties of quantum mechanics. This would, of course, mean that, in this case, if subparticles are used to obtain a cosmology, then the created cosmology would need to have some basic quantum mechanical properties. On the other hand, if subparticles are postulated, then this would not need to be the case.

6. Each of the universes has, within the NSP-world (i.e. nonstandard physical world), a beginning; even those cosmologies that claim to be cyclic. (I’ll discuss this below.)

7. The GGU-model uses only ONE structure to produce all of its effects; all of the different universes. Further, many of these universes have totally different beginnings and, yet, each would have the same appearance as viewed by any observer from a certain moment after the development

“started” and until the development “ends” in the NSP-world sense.

8. A few years ago, I separated some, but not all, of the universe creating scenarios into different categories and illustrated them in a very short paper called “Aspects of the GGU-model.” Among these scenarios are those scenarios that have portions that cannot be described by any human theory. But one of the simplest scenarios, from the NSP-world viewpoint, is the classical Big Bang. This does not mean that a humanly comprehensible description for this scenario would be “simple” for the human mind to comprehend. But it is almost trivial from the NSP-world viewpoint.

9. All scenarios that the scientific community consider, at this very moment in human history, to be “exciting” and worthy of extensive theoretical research are also almost trivial from the NSP-world viewpoint, although they may have taken great human effort to produce. This includes any of the proposed cyclic or multi-universe models.

10. Cyclic or multi-universe scenarios fit into the category I term the repeated but not necessarily identical minimum scenario. Now in the natural-world one can conjecture that such a cosmology has and will last forever, i.e. it will never cease and had no natural beginning. But in the NSP-world, it does have a beginning and will cease in a very simple way. This is due to the different NSP-world “sequence or order” properties when compared with the standard concept associated with a natural-world ordered development.

11. I’ll try not to be technical at this point, not an easy thing to do. There is no such thing in the natural-world as “infinite” time in the sense of the limit concept or say a measure with respect to the real or complex numbers. (Even if we use the extended interval, the extended plane, manifolds or whatever). All this actually means is that such a ordered measure would be unbounded (maybe in a special sense) as a real, complex, etc. number. But, such ordered measures ARE BOUNDED in the NSP-world. In fact, there is a bounded collection of ordered NSP-world measures that contains these ordered natural-world measures but behaves as if it were simply a FINITE ordered set of measures. See the discussion on page 11 of the “Solutions . . .” monograph. The results established relative to this GGU-model interpretation apply to cyclic, plasma or steady state types of “it has and will exist forever” universes. Such NSP-world scenarios are not as trivial as the Big Bang type. But they are still rather simple from the NSP-world point of view. Once again, this does not mean that a humanly comprehensible description (a cosmology) for such a scenario is easily achieved. Relative to various theories of gravitation such as the Hilbert-Einstein theory, subparticles also lead to gravitational effects, among other effects. These are discussed in the book “Einstein Corrected” where it is shown that such effects can be predicted by means of a very simple electromagnetic interaction between the substratum and the natural-world. This tends to substantiate the Poincaré statement that everything within the natural universe is related to electromagnetic properties.

12. Having more than one secular cosmological model from which to choose, even if they cannot be differentiated one from another by any form of observation that can be made today, is a good thing to have for it then requires an individual’s choice to be based upon identifiable philosophical concepts exterior to the scientific method.

13. One of the major misunderstandings is the relationship between the methods used to establish the GGU-model, modern science and scientific descriptions. Assuming that an individual uses any acceptable form of human thought and describes the behavior of a developing natural system using any form of natural science terminology, then an ultralogic restricted to the language of this scientific description will also give the exact same description for the behavior of the same developing natural system. Scientifically, why is this always the case? Well, the GGU-model has as its foundations the GENERAL properties used by the scientist to create just such a DESCRIPTION

for the behavior of a developing natural system. Also, the GGU-model is based upon the simplest form of human thought that must be used to be consider as human.

14. I repeat that the GGU-model satisfies ALL of the requirements for a pre-geometry, and it answers all of the questions. But, some scientists dislike the GGU-model for an entirely different reason – a personal and philosophical reason. The GGU-model specifically predicts that there can be no human language that can be used to give a detailed description for certain processes and events that occur within the substratum. It is this barrier to human knowledge that bothers those individuals who consider humankind as capable of knowing everything about “all there is, or ever was, or ever will be.” One the other hand, the GGU-model substantiates the philosophy of science of Max Planck who wrote, “Nature does not allow herself to be exhaustively expressed in human thought.”

15. Finally, is it possible for the NSP-world background universe and its submodel the GGU-model exist in objective reality? Since 99% of all of the mathematical analysis used within the physical sciences requires the NSP-world as a substratum, the answer is yes. In particle physics and elsewhere, the concept of predictability and the indirect verification of natural-system behavior is the concept used to assert that certain microworld entities exist in objective reality. This appears to be necessary since only macroscopic or large scale behavior can be observed either by human or machine sensors. Using this exact same philosophy of science, one would immediately conclude that the NSP-world and the GGU-model processes exist in object reality.

Sincerely,

Prof. Robert A. Herrmann